


# A new class of metric $f$ -manifolds

PABLO ALEGRE<sup>1</sup>, LUIS M. FERNÁNDEZ<sup>2</sup> and ALICIA PRIETO-MARTÍN<sup>2</sup>


**ABSTRACT.** We introduce a new general class of metric  $f$ -manifolds which we call (almost) trans- $S$ -manifolds and includes  $S$ -manifolds,  $C$ -manifolds,  $s$ -th Sasakian manifolds and generalized Kenmotsu manifolds studied previously. We prove their main properties and we present many examples which justify their study.

## 1. INTRODUCTION

In complex geometry, the relationships between the different classes of manifolds can be summarized in the well known diagram by Blair [3]:



In the case of contact geometry we have the diagram:



In the above diagram the almost contact structure  $(\phi, \eta, \xi)$  is said to be normal if  $[\phi, \phi] + 2d\eta \otimes \xi = 0$  and condition (1) is

$$(\nabla_X \phi)Y = g(X, Y)\xi - \eta(Y)X,$$

for any tangent vector fields  $X$  and  $Y$ .

Moreover, an almost contact metric manifold  $(M, \phi, \xi, \eta, g)$  is said to have an  $(\alpha, \beta)$  trans-Sasakian structure if (see [11] for more details)

$$(1.1) \quad (\nabla_X \phi)Y = \alpha\{g(X, Y)\xi - \eta(Y)X\} + \beta\{g(\phi X, Y)\xi - \eta(Y)\phi X\},$$


Received: 15.01.2018. In revised form: 30.05.2018. Accepted: 06.06.2018

2010 Mathematics Subject Classification. 53C15, 53C25, 53C99.

Key words and phrases. Almost trans- $S$ -manifold, trans- $S$ -manifold, generalized  $D$ -conformal deformation, warped product.

Corresponding author: Luis M. Fernández; [lmfer@us.es](mailto:lmfer@us.es)

where  $\alpha, \beta$  are differentiable functions (called characteristic functions) on  $M$ . Particular cases of trans-Sasakian manifolds are Sasakian ( $\alpha = 1, \beta = 0$ ), cosymplectic ( $\alpha = \beta = 0$ ) or Kenmotsu ( $\alpha = 0, \beta = 1$ ) manifolds. In fact, we can extend the above diagram to



where

$$(1.2) \quad d\Phi = \Phi \wedge (\phi^*(\delta\Phi) - (\delta\eta)\eta), \quad d\eta = \frac{1}{2n} \{ \delta\Phi(\xi)\Phi - 2\eta \wedge \phi^*(\delta\Phi) \}.$$

and:

$$(1.3) \quad d\Phi = \frac{-1}{n} \delta\eta(\Phi \wedge \eta), \quad d\eta = \frac{1}{2n} \delta\Phi(\xi)\Phi, \quad \phi^*(\delta\Phi) = 0.$$

More generally, K. Yano [15] introduced the notion of  $f$ -structure on a  $(2n + s)$ -dimensional manifold as a tensor field  $f$  of type (1,1) and rank  $2n$  satisfying  $f^3 + f = 0$ . Almost complex ( $s = 0$ ) and almost contact ( $s = 1$ ) structures are well-known examples of  $f$ -structures. In this context, D. E. Blair [2] defined  $K$ -manifolds (and particular cases of  $S$ -manifolds and  $C$ -manifolds). Then,  $K$ -manifolds are the analogue of Kaehlerian manifolds in the almost complex geometry and  $S$ -manifolds (resp.,  $C$ -manifolds) of Sasakian manifolds (resp., cosymplectic manifolds) in the almost contact geometry. Consequently, one can obtain a similar diagram for metric  $f$ -manifolds, that is, manifolds endowed with an  $f$ -structure and a compatible metric.

The purpose of the present paper is to introduce a new class of metric  $f$ -manifolds which generalizes the one of trans-Sasakian manifolds. In this context, we notice that there has been a previous generalization of  $(\alpha, 0)$ -trans-Sasakian manifolds for metric  $f$ -manifolds. It was due to I. Hasegawa, Y. Okuyama and T. Abe who introduced the so-called *homothetic  $s$ -contact Riemannian manifolds* in [8] as metric  $f$ -manifolds such that  $2c_i g(fX, Y) = d\eta_i(X, Y)$  for certain nonzero constants  $c_i, i = 1, \dots, s$  (actually, they use  $p$  instead of  $s$ ). In particular, if the structure vector fields  $\xi_i$  are Killing vector fields and the  $f$ -structure is also normal, the manifold is called a *homothetic  $s$ -th Sasakian manifold*. They proved that a homothetic  $s$ -contact Riemannian manifold is a homothetic  $s$ -th Sasakian manifold if and only if

$$(\nabla_X f)Y = - \sum_{i=1}^s c_i \{ g(fX, fY)\xi_i + \eta_i(Y)f^2X \},$$

and

$$\nabla_X \xi_i = c_i fX,$$

for any tangent vector fields  $X$  and  $Y$  and any  $i = 1, \dots, s$ .

More recently, M. Falcitelli and A. M. Pastore have introduced  $f$ -structures of Kenmotsu type as those normal  $f$ -manifolds with  $dF = 2\eta^1 \wedge F$  and  $d\eta^i = 0$  for  $i = 1, \dots, s$  [5]. In this context, L. Bhatt and K. K. Dube [1] and A. Turgut Vanli and R. Sari [14] have studied a more general type of Kenmotsu  $f$ -manifolds for which all the structure 1-forms

$\eta_i$  are closed and:

$$dF = 2 \sum_{i=1}^s \eta_i \wedge F.$$

These examples motivate the idea of introducing the mentioned new more general class of metric  $f$ -manifolds, including the above ones, which we shall call *trans-S-manifolds* because trans-Sasakian manifolds become to be a particular case of them.

## 2. METRIC $f$ -MANIFOLDS

A  $(2n + s)$ -dimensional Riemannian manifold  $(M, g)$  endowed with an  $f$ -structure  $f$  (that is, a tensor field of type  $(1,1)$  and rank  $2n$  satisfying  $f^3 + f = 0$  [15]) is said to be a *metric  $f$ -manifold* if, moreover, there exist  $s$  global vector fields  $\xi_1, \dots, \xi_s$  on  $M$  (called *structure vector fields*) such that, if  $\eta_1, \dots, \eta_s$  are the dual 1-forms of  $\xi_1, \dots, \xi_s$ , then

$$(2.4) \quad \begin{aligned} f\xi_\alpha &= 0; \eta_\alpha \circ f = 0; f^2 = -I + \sum_{\alpha=1}^s \eta_\alpha \otimes \xi_\alpha; \\ g(X, Y) &= g(fX, fY) + \sum_{i=1}^s \eta_i(X)\eta_i(Y), \end{aligned}$$

for any  $X, Y \in \mathcal{X}(M)$  and  $i = 1, \dots, s$ . The distribution on  $M$  spanned by the structure vector fields is denoted by  $\mathcal{M}$  and its complementary orthogonal distribution is denoted by  $\mathcal{L}$ . Consequently,  $TM = \mathcal{L} \oplus \mathcal{M}$ . Moreover, if  $X \in \mathcal{L}$ , then  $\eta_\alpha(X) = 0$ , for any  $\alpha = 1, \dots, s$  and if  $X \in \mathcal{M}$ , then  $fX = 0$ .

For a metric  $f$ -manifold  $M$  we can construct very useful local orthonormal basis of tangent vector fields. To this end, let  $U$  be a coordinate neighborhood on  $M$  and  $X_1$  any unit vector field on  $U$ , orthogonal to the structure vector fields. Then,  $fX_1$  is another unit vector field orthogonal to  $X_1$  and to the structure vector fields too. Now, if it is possible, we choose a unit vector field  $X_2$  orthogonal to the structure vector fields, to  $X_1$  and to  $fX_1$ . Then,  $fX_2$  is also a unit vector field orthogonal to the structure vector fields, to  $X_1$ , to  $fX_1$  and to  $X_2$ . Proceeding in this way, we obtain a local orthonormal basis  $\{X_i, fX_i, \xi_j\}$ ,  $i = 1, \dots, n$  and  $j = 1, \dots, s$ , called an  *$f$ -basis*.

Let  $F$  be the 2-form on  $M$  defined by  $F(X, Y) = g(X, fY)$ , for any  $X, Y \in \mathcal{X}(M)$ . Since  $f$  is of rank  $2n$ , then

$$\eta_1 \wedge \dots \wedge \eta_s \wedge F^n \neq 0$$

and, in particular,  $M$  is orientable. A metric  $f$ -manifold is said to be a *metric  $f$ -contact manifold* if  $F = d\eta_i$ , for any  $i = 1, \dots, s$ .

The  $f$ -structure  $f$  is said to be *normal* if

$$[f, f] + 2 \sum_{i=1}^s \xi_i \otimes d\eta_i = 0,$$

where  $[f, f]$  denotes the Nijenhuis tensor of  $f$ . If  $f$  is normal, then [7]

$$(2.5) \quad [\xi_i, \xi_j] = 0,$$

for any  $i, j = 1, \dots, s$ .

A metric  $f$ -manifold is said to be a  *$K$ -manifold* [2] if it is normal and  $dF = 0$ . In a  $K$ -manifold  $M$ , the structure vector fields are Killing vector fields [2]. A  $K$ -manifold is called an  *$S$ -manifold* if  $F = d\eta_i$ , for any  $i$  and a  *$C$ -manifold* if  $d\eta_i = 0$ , for any  $i$ . Note that, for  $s = 0$ , a  $K$ -manifold is a Kaehlerian manifold and, for  $s = 1$ , a  $K$ -manifold is a quasi-Sasakian manifold, an  $S$ -manifold is a Sasakian manifold and a  $C$ -manifold

is a cosymplectic manifold. When  $s \geq 2$ , non-trivial examples can be found in [2, 8]. Moreover, a  $K$ -manifold  $M$  is an  $S$ -manifold if and only if

$$(2.6) \quad \nabla_X \xi_i = -fX, \quad X \in \mathcal{X}(M), \quad i = 1, \dots, s,$$

and it is a  $C$ -manifold if and only if

$$(2.7) \quad \nabla_X \xi_i = 0, \quad X \in \mathcal{X}(M), \quad i = 1, \dots, s.$$

It is easy to show that in an  $S$ -manifold,

$$(2.8) \quad (\nabla_X f)Y = \sum_{i=1}^s \{g(fX, fY)\xi_i + \eta_i(Y)f^2X\},$$

for any  $X, Y \in \mathcal{X}(M)$  and in a  $C$ -manifold,

$$(2.9) \quad \nabla f = 0.$$

### 3. DEFINITION OF TRANS-S-MANIFOLDS AND MAIN PROPERTIES

The original idea to define  $(\alpha, \beta)$  trans-Sasakian manifolds is to generalize cosymplectic, Kenmotsu and Sasakian manifolds.

Kenmotsu: $d\eta = 0, \text{ normal}$	Quasi-Sasakian: $d\Phi = 0,$ normal	Trans-Sasakian: $d\Phi = 2\beta(\Phi \wedge \eta),$ $d\eta = \alpha\Phi,$ $\phi^*(\delta\Phi) = 0,$ normal
Cosymplectic: $d\Phi = 0, \quad d\eta = 0,$ normal		
Sasakian: $\Phi = d\eta, \text{ normal}$		

In the same way, our idea is to define *trans-S-manifolds* generalizing  $C$ -manifolds,  $f$ -Kenmotsu and  $S$ -manifolds.

As we said in the Introduction, an almost contact manifold is trans-Sasakian if and only if it (1.1) holds. Then, we can introduce the notion of trans- $S$ -manifold in a similar way.

**Definition 3.1.** A  $(2n + s)$ -dimensional metric  $f$ -manifold  $M$  is said to be a **almost trans- $S$ -manifold** if it satisfies

$$(3.10) \quad \begin{aligned} (\nabla_X f)Y = \sum_{i=1}^s [\alpha_i \{g(fX, fY)\xi_i + \eta_i(Y)f^2X\} \\ + \beta_i \{g(fX, Y)\xi_i - \eta_i(Y)fX\}], \end{aligned}$$

for certain smooth functions (called the **characteristic functions**)  $\alpha_i, \beta_i, i = 1, \dots, s$ , on  $M$  and any  $X, Y \in \mathcal{X}(M)$ . If, moreover,  $M$  is normal, then it is said to be a **trans- $S$ -manifold**.

So, if  $s = 1$ , a trans- $S$ -manifold is actually a trans-Sasakian manifold. Furthermore, in this case, condition (3.10) implies normality. However, for  $s \geq 2$ , this does not hold. In fact, it is straightforward to prove that, for any  $X, Y \in \mathcal{X}(M)$ ,

$$(3.11) \quad [f, f](X, Y) + 2 \sum_{i=1}^s d\eta_i(X, Y)\xi_i = \sum_{i,j=1}^s [\eta_j(\nabla_X \xi_i)\eta_j(Y) - \eta_j(\nabla_Y \xi_i)\eta_j(X)] \xi_i,$$

which is not zero in general. But, in a trans- $S$ -manifold, (3.11) implies that, for any  $X \in \mathcal{X}(M)$  and any  $i = 1, \dots, s$ :

$$\sum_{j=1}^s \eta_j(\nabla_X \xi_i) \eta_j(Y) - \sum_{j=1}^s \eta_j(\nabla_Y \xi_i) \eta_j(X) = 0.$$

If we put  $Y = \xi_k$ , from (2.5) we get that

$$(3.12) \quad \eta_k(\nabla_X \xi_i) = 0,$$

for any  $i, k = 1, \dots, s$ . Using this fact, from (3.10), we deduce that

$$(3.13) \quad \nabla_X \xi_i = -\alpha_i fX - \beta_i f^2 X,$$

for any  $X \in \mathcal{X}(M)$  and any  $i = 1, \dots, s$ .

Now, we can prove:

**Theorem 3.1.** *A almost trans- $S$ -manifold  $M$  is a trans- $S$ -manifold if and only if (3.13) holds for any  $X \in \mathcal{X}(M)$  and any  $i = 1, \dots, s$ .*

*Proof.* From (3.10) we have that, for any  $X \in \mathcal{X}(M)$  and any  $i = 1, \dots, s$ :

$$\nabla_X \xi_i = -\alpha_i fX - \beta_i f^2 X + \sum_{j=1}^s \eta_j(\nabla_X \xi_i) \xi_j.$$

Comparing this equality and (3.13) we have that  $\eta_j(\nabla_X \xi_i) = 0$ , for any  $i, j = 1, \dots, s$ . So, from (3.11), the metric  $f$ -manifold  $M$  is normal and, consequently, a trans- $S$ -manifold. The converse is obvious.  $\square$

Observe that (3.10) can be re-written as

$$\begin{aligned} (\nabla_X F)(Y, Z) = & \sum_{i=1}^s [\alpha_i \{g(fX, fZ) \eta_i(Y) - g(fX, fY) \eta_i(Z)\} \\ & + \beta_i \{g(X, fY) \eta_i(Z) - g(X, fZ) \eta_i(Y)\}], \end{aligned}$$

for any  $X, Y, Z \in \mathcal{X}(M)$ . Then, if  $X \in \mathcal{L}$  is a unit vector field, we have:

$$(\nabla_X F)(X, \xi_i) = -\alpha_i, \quad (\nabla_X F)(\xi_i, fX) = \beta_i, \quad i = 1, \dots, s.$$

Moreover, from (3.13), we deduce

$$(\nabla_X \eta_i)Y = \alpha_i g(X, fY) + \beta_i g(fX, fY),$$

for any  $X, Y \in \mathcal{X}(M)$  and any  $i = 1, \dots, s$ . Again, if  $X \in \mathcal{L}$  is a unit vector field, we get:

$$(\nabla_X \eta_i)fX = -\alpha_i, \quad (\nabla_X \eta_i)X = \beta_i, \quad i = 1, \dots, s.$$

For trans- $S$ -manifolds, we can prove the following theorem.

**Theorem 3.2.** *Let  $M$  be a trans- $S$ -manifold. Then,  $(\delta F)\xi_i = 2n\alpha_i$  and  $\delta\eta_i = -2n\beta_i$ , for any  $i = 1, \dots, s$ .*

*Proof.* Taking a  $f$ -basis  $\{X_1, \dots, X_n, fX_1, \dots, fX_n, \xi_1, \dots, \xi_s\}$ , since

$$\begin{aligned} (\delta F)X = & - \sum_{k=1}^n \{(\nabla_{X_k} F)(X_k, X) + (\nabla_{fX_k} F)(fX_k, X)\} \\ & - \sum_{j=1}^s (\nabla_{\xi_j} F)(\xi_j, X) \\ = & \sum_{k=1}^n \{g(X_k, (\nabla_{X_k} \phi)X) + g(fX_k, (\nabla_{fX_k} \phi)X)\}, \end{aligned}$$

for any  $X \in \mathcal{X}(M)$ , by using (3.10) it is straightforward to obtain

$$(3.14) \quad (\delta F)X = 2n \sum_{j=1}^s \alpha_j \eta_j(X)$$

and, putting  $X = \xi_i$ , it follows that  $(\delta F)\xi_i = 2n\alpha_i$ .

Moreover,

$$\delta\eta_i = - \sum_{k=1}^n \{(\nabla_{X_k} \eta_i)X_k + (\nabla_{fX_k} \eta_i)fX_k\} - \sum_{j=1}^s (\nabla_{\xi_j} \eta_i)\xi_j,$$

for any  $i = 1, \dots, s$ . But, from (3.12) we get that  $(\nabla_{\xi_j} \eta_i)\xi_j = 0$ , for any  $j = 1, \dots, s$ . Consequently, by using (3.13)

$$\begin{aligned} \delta\eta_i &= - \sum_{k=1}^n \{g(X_k, \nabla_{X_k} \xi_i) + g(fX_k, \nabla_{fX_k} \xi_i)\} \\ &= - \sum_{k=1}^n \beta_i \{g(X_k, X_k) + g(fX_k, fX_k)\} = -2n\beta_i. \end{aligned}$$

which concludes the proof.  $\square$

The above theorem generalizes the result given by D. E. Blair and J. A. Oubiña in [4] for trans-Sasakian manifolds. Moreover, trans- $S$ -manifolds verify certain desirable conditions.

**Proposition 3.1.** *Let  $M$  be a trans- $S$ -manifold. The following equations are verified:*

- (i)  $dF = 2F \wedge \sum_{i=1}^s \beta_i \eta_i$ ;
- (ii)  $d\eta_i = \alpha_i F, i = 1, \dots, s$ ;
- (iii)  $f^*(\delta F) = 0$ .

*Proof.* From (3.10), a direct computation gives, for any  $X, Y, Z \in \mathcal{X}(M)$ :

$$\begin{aligned} dF(X, Y, Z) &= -g((\nabla_X f)Y, Z) + g((\nabla_Y f)X, Z) - g((\nabla_Z f)X, Y) \\ &= 2 \sum_{i=1}^s \{-\beta_i \eta_i(Z)g(fX, Y) + \beta_i \eta_i(Y)g(fX, Z) - \beta_i \eta_i(X)g(fY, Z)\} \\ &= 2(F \wedge \sum_{i=1}^s \beta_i \eta_i)(X, Y, Z). \end{aligned}$$

Next, from (3.13) it is obtained the second statement. Finally, from (3.14) we get (iii).  $\square$

From (ii) of the above proposition we observe that if one of the functions  $\alpha_i$  is a non-zero constant function, then the 2-form  $F$  is closed and the trans- $S$ -manifold  $M$  is a  $K$ -manifold. Moreover we can prove:

**Theorem 3.3.** *A trans- $S$ -manifold  $M$  is a  $K$ -manifold if and only if  $\beta_1 = \dots = \beta_s = 0$ .*

*Proof.* Firstly, if all the functions  $\beta_i$  are equal to zero, from (i) of Proposition 3.1 we get  $dF = 0$  and  $M$  is a  $K$ -manifold.

Conversely, it is known (see [6]) that, for  $K$ -manifolds, the following formula holds, for any  $X, Y, Z \in \mathcal{X}(M)$ :

$$g((\nabla_X f)Y, Z) = \sum_{i=1}^s \{d\eta_i(fY, X)\eta_i(Z) - d\eta_i(fZ, X)\eta_i(Y)\}.$$

Consequently, from (ii) of Proposition 3.1 and (3.10) we conclude that  $\beta_i = 0$ , for any  $i = 1, \dots, s$ .  $\square$

From Theorem 3.2 we deduce:

**Corollary 3.1.** *A trans- $S$ -manifold  $M$  is a  $K$ -manifold if and only if  $\delta\eta_i = 0$ , for any  $i = 1, \dots, s$ .*

Furthermore, taking into account (2.6) and (2.7), we have:

**Corollary 3.2.** *Any trans- $S$ -manifold is an  $S$ -manifold if and only if  $\alpha_i = 1, \beta_i = 0$  and it is a  $C$ -manifold if and only if  $\alpha_i = \beta_i = 0$ , in both cases for any  $i = 1, \dots, s$ .*

In next section, we shall present some examples of trans- $S$ -manifolds which are not  $K$ -manifolds due to the fact that not all their characteristic functions  $\beta_i$  are zero. Now, the natural question is if any  $K$ -manifold is a trans- $S$ -manifold. In general, the answer is negative and to this end, we can consider the following example.

Let  $(N, J, G)$  be a Kaehler manifold,  $(M, f, \xi_1, \dots, \xi_s, \eta_1, \dots, \eta_s, g)$  be an  $S$ -manifold and  $\widetilde{M} = N \times M$ .

If  $\widetilde{X} = U + X, \widetilde{Y} = V + Y \in \mathcal{X}(\widetilde{M})$ , where  $U, V \in \mathcal{X}(N)$  and  $X, Y \in \mathcal{X}(M)$ , respectively, we can define a metric  $f$ -structure on  $\widetilde{M}$  by the following structure elements:

$$\begin{aligned} \widetilde{f}(U + X) &= JU + fX, \quad \widetilde{\xi}_i = 0 + \xi_i, \quad \widetilde{\eta}_i(U + X) = \eta_i(X), \quad i = 1, \dots, s, \\ \widetilde{g}(U + X, V + Y) &= G(U, V) + g(X, Y). \end{aligned}$$

It is straightforward to check that  $\widetilde{M}$  with this structure is a  $K$ -manifold. However, it is not a trans- $S$ -manifold. In fact, since  $N$  is a Kaehler manifold and so,  $J$  is parallel, if  $\nabla$  and  $\widetilde{\nabla}$  denote the Riemannian connections of  $M$  and  $\widetilde{M}$ , respectively, then

$$(\widetilde{\nabla}_{\widetilde{X}} \widetilde{f})\widetilde{Y} = 0 + (\nabla_X f)Y$$

and, consequently, (3.10) does not hold for  $\widetilde{M}$ .

However, we can observe that, from (2.8) and (2.9), the particular cases of  $S$ -manifolds and  $C$ -manifolds are trans- $S$ -manifolds.

On the other hand, it is known [2] that, in a  $K$ -manifold, all the structure vector fields are Killing vector fields. For trans- $S$ -manifolds we can prove:

**Proposition 3.2.** *Let  $M$  be a trans- $S$ -manifold. Then, the structure vector field  $\xi_i$  is a Killing vector field if and only if the corresponding characteristic function  $\beta_i = 0$ .*

*Proof.* A direct computation by using (3.13) gives

$$(L_{\xi_i} g)(X, Y) = 2\beta_i g(fX, fY),$$

for any  $X, Y \in \mathcal{X}(M)$ . This completes the proof.  $\square$

#### 4. EXAMPLES OF TRANS- $S$ -MANIFOLDS

As we have mentioned above, it is obvious that, from (2.8) and (2.9),  $S$ -manifolds and  $C$ -manifolds are trans- $S$ -manifolds. Moreover, the homothetic  $s$ -th Sasakian manifolds of [8] are also trans- $S$ -manifolds with the function  $\alpha_i$  constant and  $\beta_i = 0$ , for any  $i$ .

From Propositions 2.2 and 2.5 of [5], we see that  $f$ -manifolds of Kenmotsu type, introduced by M. Falcitelli and A.M. Pastore, actually are trans- $S$ -manifolds with functions  $\alpha_1 = \dots = \alpha_s = \beta_2 = \dots = \beta_s = 0$  and  $\beta_1 = 1$ .

Also, from Theorem 2.4 in [14], we see that generalized Kenmotsu manifolds studied by L. Bhatt and K.K. Dube [1] and A. Turgut Vanli and R. Sari [14] are trans- $S$ -manifolds with functions  $\alpha_1 = \dots = \alpha_s = 0$  and  $\beta_1 = \dots = \beta_s = 1$ .

Then, we are going to look for examples with different non-constant functions  $\alpha_i$  and  $\beta_i$ . We shall obtain these examples by using  $D$ -conformal deformations and warped products.

Firstly, given a metric  $f$ -manifold  $(M, f, \xi_1, \dots, \xi_s, \eta_1, \dots, \eta_s, g)$ , let us consider the generalized  $D$ -conformal deformation given by

$$(4.15) \quad \tilde{f} = f, \quad \tilde{\xi}_i = \frac{1}{a}\xi_i, \quad \tilde{\eta}_i = a\eta_i, \quad \tilde{g} = bg + (a^2 - b) \sum_{i=1}^s \eta_i \otimes \eta_i,$$

for any  $i = 1, \dots, s$ , where  $a, b$  are two positive differentiable functions on  $M$ . Then, it is easy to see that  $(M, \tilde{f}, \tilde{\xi}_1, \dots, \tilde{\xi}_s, \tilde{\eta}_1, \dots, \tilde{\eta}_s, \tilde{g})$  is also a metric  $f$ -manifold. Let us notice that we can obtain conformal,  $D$ -homothetic (see [13]) or  $D$ -conformal (in the sense of S. Suguri and S. Nakayama [12]) deformations, by putting  $a^2 = b$ ,  $a = b = \text{constant}$  or  $a = b$  in (4.15), respectively. In [9] Z. Olszack considered  $a$  and  $b$  constants,  $a \neq 0$ ,  $b > 0$  but not necessarily equal and he also called the resulting transformation a  $D$ -homothetic deformation.

Moreover, let us suppose that  $M$  is a trans- $S$ -manifold and that  $a, b$  depend only on the directions of the structure vector fields  $\xi_i$ ,  $i = 1, \dots, s$ . Therefore, we can calculate  $\tilde{\nabla}$  from  $\nabla$  and  $\tilde{g}$  by using Koszul's formula and (3.13). It follows that the Riemannian connection  $\tilde{\nabla}$  of  $\tilde{g}$  is given by

$$(4.16) \quad \begin{aligned} \tilde{\nabla}_X Y &= \nabla_X Y + \sum_{i=1}^s \frac{2(a^2 - b)\beta_i - \xi_i b}{2a^2} g(fX, fY) \xi_i \\ &\quad - \frac{1}{2b} \{ (Xb)f^2 Y + (Yb)f^2 X \} \\ &\quad + \frac{1}{2a^2} \sum_{i=1}^s \{ (Xa^2)\eta_i(Y) + (Ya^2)\eta_i(X) \\ &\quad - (\xi_i a^2) \sum_{j=1}^s \eta_j(X)\eta_j(Y) \} \xi_i \\ &\quad - \frac{a^2 - b}{b} \sum_{i=1}^s \alpha_i \{ \eta_i(Y)fX + \eta_i(X)fY \}, \end{aligned}$$

for any vector fields  $X, Y \in \mathcal{X}(M)$ .

**Theorem 4.4.** *Let  $(M, f, \xi_1, \dots, \xi_s, \eta_1, \dots, \eta_s, g)$  be a trans- $S$ -manifold and consider a generalized  $D$ -conformal deformation on  $M$ , with  $a, b$  positive functions depending only on the directions of the structure vector fields. Then  $(M, \tilde{f}, \tilde{\xi}_1, \dots, \tilde{\xi}_s, \tilde{\eta}_1, \dots, \tilde{\eta}_s, \tilde{g})$  is also a trans- $S$ -manifold with functions:*

$$\tilde{\alpha}_i = \frac{\alpha_i a}{b}, \quad \tilde{\beta}_i = \frac{\xi_i b}{2ab} + \frac{\beta_i}{a}, \quad i = 1, \dots, s.$$

*Proof.* By using (4.16) and taking into account that  $b$  only depends on the directions of the structure vector fields, we have

$$\begin{aligned} (\tilde{\nabla}_X \tilde{f})Y &= (\nabla_X f)Y - \sum_{i=1}^s \frac{2(a^2 - b)\beta_i - \xi_i b}{2a^2} g(fX, Y) \xi_i \\ &\quad - \frac{1}{2b} \sum_{i=1}^s (\xi_i b) \eta_i(Y) fX + \frac{a^2 - b}{b} \sum_{i=1}^s \alpha_i \eta_i(Y) f^2 X, \end{aligned}$$



for any  $X, Y \in \mathcal{X}(M)$ . Now, since  $M$  is trans- $S$ -manifold, from (3.10) and (4.15) we obtain

$$(\tilde{\nabla}_X \tilde{f})Y = \sum_{i=1}^s \left\{ \frac{\alpha_i a}{b} (\tilde{g}(\tilde{f}X, \tilde{f}Y)\tilde{\xi}_i + \tilde{\eta}_i(Y)X) + \left( \frac{\xi_i b}{2ab} + \frac{\beta_i}{a} \right) (\tilde{g}(\tilde{f}X, Y)\tilde{\xi}_i - \tilde{\eta}_i(Y)\tilde{f}X) \right\},$$

and this completes the proof.  $\square$

Note that if  $M$  is a Sasakian manifold, that is, if  $s = 1, \alpha = 1$  and  $\beta = 0$ , this method does not produce an  $(\alpha, \beta)$  trans-Sasakian manifold but an  $(\alpha, 0)$  one because, by Darboux's theorem, if  $a, b$  only depend of the direction of  $\xi$ , they should be constants.

**Corollary 4.3.** *Let  $(M, f, \xi_1, \dots, \xi_s, \eta_1, \dots, \eta_s, g)$  be an  $S$ -manifold and consider a generalized  $D$ -conformal deformation on  $M$ , with  $a, b$  positive functions depending only on the directions of the structure vector fields. Then  $(M, \tilde{f}, \tilde{\xi}_1, \dots, \tilde{\xi}_s, \tilde{\eta}_1, \dots, \tilde{\eta}_s, \tilde{g})$  is a trans- $S$ -manifold with functions:*

$$\tilde{\alpha}_i = \frac{a}{b}, \quad \tilde{\beta}_i = \frac{\xi_i b}{2ab}, \quad i = 1, \dots, s.$$

**Corollary 4.4.** *Let  $(M, f, \xi_1, \dots, \xi_s, \eta_1, \dots, \eta_s, g)$  be an  $C$ -manifold and consider a generalized  $D$ -conformal deformation on  $M$ , with  $a, b$  positive functions depending only on the directions of the structure vector fields. Then  $(M, \tilde{f}, \tilde{\xi}_1, \dots, \tilde{\xi}_s, \tilde{\eta}_1, \dots, \tilde{\eta}_s, \tilde{g})$  is a trans- $S$ -manifold with functions:*

$$\tilde{\alpha}_i = 0, \quad \tilde{\beta}_i = \frac{\xi_i b}{2ab}, \quad i = 1, \dots, s.$$

**Corollary 4.5.** *Let  $(M, f, \xi_1, \dots, \xi_s, \eta_1, \dots, \eta_s, g)$  be a generalized Kenmotsu manifold and consider a generalized  $D$ -conformal deformation on  $M$ , with  $a, b$  positive functions depending only on the directions of the structure vector fields. Then  $(M, \tilde{f}, \tilde{\xi}_1, \dots, \tilde{\xi}_s, \tilde{\eta}_1, \dots, \tilde{\eta}_s, \tilde{g})$  is a trans- $S$ -manifold with functions:*

$$\tilde{\alpha}_i = 0, \quad \tilde{\beta}_i = \frac{\xi_i b}{2ab} + \frac{1}{a}, \quad i = 1, \dots, s.$$

Next, we are going to construct more examples of trans- $S$ -manifolds by using warped products. For later use, we need the following lemma from [10] to compute the Riemannian connection of a warped product:

**Lemma 4.1.** *Let us consider  $M = B \times_h F$  and denote by  $\nabla, \nabla^B$  and  $\nabla^F$  the Riemannian connections on  $M, B$  and  $F$ . If  $X, Y$  are tangent vector fields on  $B$  and  $V, W$  are tangent vector fields on  $F$ , then:*

- (i)  $\nabla_X Y$  is the lift of  $\nabla_X^B Y$ .
- (ii)  $\nabla_X V = \nabla_V X = (Xh/h)V$ .
- (iii) The component of  $\nabla_V W$  normal to the fibers is:

$$-(g_h(V, W)/h)\text{grad } h.$$

- (iv) The component of  $\nabla_V W$  tangent to the fibers is the lift of  $\nabla_V^F W$ .

In this context, if  $(N, J, G)$  is an almost Hermitian manifold, the warped product  $\tilde{M} = \mathbb{R}^s \times_h N$  can be endowed with a metric  $f$ -structure

$$(\tilde{f}, \tilde{\xi}_1, \dots, \tilde{\xi}_s, \tilde{\eta}_1, \dots, \tilde{\eta}_s, g_h),$$

with the warped metric  $g_h = -\pi^*(g_{\mathbb{R}^s}) + (h \circ \pi)^2 \sigma^*(G)$ , where  $h > 0$  is a differentiable function on  $\mathbb{R}^s$  and  $\pi$  and  $\sigma$  are the projections from  $\mathbb{R}^s \times N$  on  $\mathbb{R}^s$  and  $N$ , respectively. In fact,  $\tilde{f}(\tilde{X}) = (J\sigma_*\tilde{X})^*$ , for any vector field  $\tilde{X} \in \mathcal{X}(\tilde{M})$  and  $\tilde{\xi}_i = \partial/\partial t_i, i = 1, \dots, s$ , where  $t_i$  denotes the coordinates of  $\mathbb{R}^s$ . Note that this metric is the one used to construct the Robertson-Walker spaces (see [10]).

Now, we study the structure of this warped product.

**Theorem 4.5.** *Let  $N$  be an almost Hermitian manifold. Then, the warped product  $(\widetilde{M} = \mathbb{R}^s \times_h N, \widetilde{f}, \widetilde{\xi}_1, \dots, \widetilde{\xi}_s, \widetilde{\eta}_1, \dots, \widetilde{\eta}_s, g_h)$  is a trans- $S$ -manifold with functions  $\widetilde{\alpha}_1 = \dots = \widetilde{\alpha}_s = 0$  and  $\widetilde{\beta}_i = h^i/h, i = 1, \dots, s$ , if and only if  $N$  is a Kaehlerian manifold, where  $h^i$  are denoting the components of the gradient of the function  $h$ , for  $i = 1, \dots, s$ .*

*Proof.* Consider  $\widetilde{X} = U + X$  and  $\widetilde{Y} = V + Y$ , where  $U, V$  and  $X, Y$  are tangent vector fields on  $\mathbb{R}^s$  and  $N$ , respectively. Then, taking into account Lemma 4.1, if  $\widetilde{\nabla}$  and  $\nabla^N$  denote the Riemannian connections of  $\widetilde{M}$  and  $N$ , respectively, we have:

$$\begin{aligned} (\widetilde{\nabla}_{\widetilde{X}} \widetilde{f}) \widetilde{Y} &= \widetilde{\nabla}_U JX + \nabla_X JY \\ &\quad - \widetilde{f}(\widetilde{\nabla}_U V + \widetilde{\nabla}_X V + \widetilde{\nabla}_U Y + \widetilde{\nabla}_X Y) \\ &= \frac{U(h)}{h} JY - \frac{g_h(X, JY)}{h} \text{grad}(h) + \nabla_X^N JY \\ &\quad - f(\nabla_U V + \frac{V(h)}{h} X + \frac{U(h)}{h} Y - \frac{g_h(X, Y)}{h} \text{grad}(h) + \nabla_X Y) \\ &= -\frac{g_h(X, JY)}{h} \text{grad}(h) - \frac{V(h)}{h} JX + (\nabla_X^N J)Y \\ &= \frac{g_h(JX, Y)}{h} \sum_{i=1}^s h^i \widetilde{\xi}_i - \sum_{i=1}^s \widetilde{\eta}_i(V) \frac{h^i}{h} JX + (\nabla_X^N J)Y \\ &= \frac{g_h(\widetilde{f}\widetilde{X}, \widetilde{Y})}{h} \sum_{i=1}^s h^i \widetilde{\xi}_i - \sum_{i=1}^s \widetilde{\eta}_i(V) \frac{h^i}{h} \widetilde{f}\widetilde{X} + (\nabla_X^N J)Y. \end{aligned}$$

Therefore, (3.10) holds if and only if  $(\nabla_U^N J)V = 0$ , that is, if and only if  $N$  is a Kaehlerian manifold. Moreover, for any  $i = 1, \dots, s$ ,

$$\widetilde{\nabla}_{\widetilde{X}} \widetilde{\xi}_i = \nabla_U \widetilde{\xi}_i + \nabla_X \widetilde{\xi}_i = \frac{h^i}{h} X = \frac{h^i}{h} (\widetilde{X} - \sum_{i=1}^s \widetilde{\eta}_i(\widetilde{X}) \widetilde{\xi}_i) = -\frac{h^i}{h} \widetilde{f}^2 \widetilde{X}$$

and then, Theorem 3.1 gives the result. □

**Corollary 4.6.** *The warped product  $\mathbb{R}^s \times_h N$ , being  $N$  a Kaehlerian manifold and  $h$  a constant function, is a  $C$ -manifold. In particular, if  $h = 1$ , the Riemannian product  $\mathbb{R}^s \times N$  is a  $C$ -manifold.*

Combining these examples with a generalized  $D$ -conformal deformation, a great variety of non-trivial trans- $S$ -manifolds can be presented.

Moreover, if we do the warped product of  $\mathbb{R}^s$  with a  $(2n + s_1)$ -dimensional (almost) trans- $S$ -manifold  $(M, f, \xi_1, \dots, \xi_{s_1}, \eta_1, \dots, \eta_{s_1}, g)$ , we obtain a new metric  $f$ -manifold

$$(\widetilde{M} = \mathbb{R}^s \times_h M, \widetilde{f}, \widetilde{\xi}_1, \dots, \widetilde{\xi}_{s+s_1}, \widetilde{\eta}_1, \dots, \widetilde{\eta}_{s+s_1}, g_h),$$

where  $\widetilde{f}(\widetilde{X}) = (f\sigma_*\widetilde{X})^*$  and:

$$\widetilde{\xi}_i = \begin{cases} \frac{\partial}{\partial t_i} & \text{if } 1 \leq i \leq s, \\ \frac{1}{h} \xi_{i-s} & \text{if } s+1 \leq i \leq s+s_1. \end{cases}$$

These manifolds, under certain hypothesis about the function  $h$ , verify (3.10) but not (3.13), so from Theorem 3.1 they are not normal. Consequently, they are examples of almost trans- $S$ -manifolds not trans- $S$ -manifolds.

**Theorem 4.6.** *Let  $M$  be a  $(2n + s_1)$ -dimensional (almost) trans- $S$ -manifold with characteristic functions  $(\alpha_i, \beta_i)$ ,  $i = 1, \dots, s_1$ . Then, the warped product  $\widetilde{M} = \mathbb{R}^s \times_h M$ , with the metric  $f$ -structure defined above, is a  $(2n + s + s_1)$ -dimensional almost trans- $S$ -manifold with functions*

$$\widetilde{\alpha}_i = \begin{cases} 0 & \text{for } i = 1, \dots, s, \\ \frac{\alpha_{i-s}}{h} & \text{for } i = s + 1, \dots, s + s_1. \end{cases}$$

and:

$$\widetilde{\beta}_i = \begin{cases} \frac{h^i}{h} & \text{for } i = 1, \dots, s, \\ \frac{\beta_{i-s}}{h} & \text{for } i = s + 1, \dots, s + s_1. \end{cases}$$

*Proof.* Consider  $\widetilde{X} = U + X$  and  $\widetilde{Y} = V + Y$ , where  $U, V$  and  $X, Y$  are tangent vector fields on  $\mathbb{R}^s$  and  $M$ , respectively. Then, taking into account Lemma 4.1, if  $\nabla$  is the Riemannian connection of  $M$ , we deduce:

$$\begin{aligned} (\nabla_{\widetilde{X}} \widetilde{f}) \widetilde{Y} &= -\frac{g_h(X, fY)}{h} \text{grad}(h) - \frac{V(h)}{h} fX + (\nabla_X f)Y \\ &= \frac{g_h(fX, Y)}{h} \sum_{i=1}^s h^i \frac{\partial}{\partial t_i} - \sum_{i=1}^s V(t_i) \frac{h^i}{h} fX \\ &\quad + \sum_{i=s+1}^{s+s_1} [\alpha_{i-s} \{g(fX, fY)\xi_{i-s} + \eta_{i-s}(Y)f^2X\} \\ &\quad + \beta_{i-s} \{g(fX, Y)\xi_{i-s} - \eta_{i-s}(Y)fX\}] \\ &= \sum_{i=1}^s \frac{h^i}{h} \{g_h(\widetilde{f}\widetilde{X}, \widetilde{Y})\xi_i - \widetilde{\eta}_i(\widetilde{Y})\widetilde{f}\widetilde{X}\} \\ &\quad + \sum_{i=s+1}^{s+s_1} \left[ \frac{\alpha_{i-s}}{h} \{g_h(\widetilde{f}\widetilde{X}, \widetilde{f}\widetilde{Y})\xi_{i-s} + \widetilde{\eta}_{i-s}(\widetilde{Y})\widetilde{f}^2\widetilde{X}\} \right. \\ &\quad \left. + \frac{\beta_{i-s}}{h} \{g_h(\widetilde{f}\widetilde{X}, \widetilde{Y})\xi_{i-s} - \widetilde{\eta}_{i-s}(\widetilde{Y})\widetilde{f}\widetilde{X}\} \right]. \end{aligned}$$

Joining the addends appropriately, it takes the form of (3.10) with the desired functions. Therefore,  $\widetilde{M}$  is a almost trans- $S$ -manifold.  $\square$

Observe that, in the above conditions, (3.13) is not verified in general. In fact, consider  $\widetilde{\xi}_i$  with  $1 \leq i \leq s$ . Then, for any  $\widetilde{X} \in \mathcal{X}(\widetilde{M})$ ,

$$\widetilde{\nabla}_{\widetilde{X}} \widetilde{\xi}_i = \frac{h^i}{h} U = \frac{h^i}{h} (\widetilde{X} - \sum_{j=1}^s \widetilde{\eta}_j(\widetilde{X})\xi_j)$$

and so, if  $h$  is not a constant function, from Theorem 3.1, we get that  $\widetilde{M}$  is not a trans- $S$ -manifold.

**Corollary 4.7.** *The warped product  $\mathbb{R}^s \times_h M$ , being  $M$  a trans- $S$ -manifold, is a trans- $S$ -manifold if and only if  $h$  is constant. In particular, the Riemannian product  $\mathbb{R}^s \times M$  is a trans- $S$ -manifold with functions*

$$(0, \cdot^s, 0, \alpha_1, \dots, \alpha_{s_1}, 0, \cdot^s, 0, \beta_1, \dots, \beta_{s_1}),$$

where  $(\alpha_i, \beta_i)$ ,  $i = 1, \dots, s_1$ , denote the characteristic functions of  $M$ .

**Corollary 4.8.** *Let  $M$  be a Sasakian manifold. Then, the warped product  $\mathbb{R} \times_h M$  is a almost trans- $S$ -manifold with functions:*

$$\alpha_1 = 0, \alpha_2 = \frac{1}{h}, \beta_1 = \frac{h'}{h} \text{ and } \beta_2 = 0.$$

**Corollary 4.9.** *Let  $M$  be a three dimensional trans-Sasakian, with non-constant characteristic functions  $\alpha$  and  $\beta$ . Then, the warped product  $\mathbb{R} \times_h M$  is a four dimensional almost trans- $S$ -manifold not trans- $S$ -manifold with functions:*

$$\alpha_1 = 0, \alpha_2 = \frac{\alpha}{h}, \beta_1 = \frac{h'}{h} \text{ and } \beta_2 = \frac{\beta}{h}.$$

## REFERENCES

- [1] Bhatt, L. and Dube, K. K., *Semi-invariant submanifolds of  $r$ -Kenmotsu manifolds*, Acta Cienc. Indica Math., **29** (2003), No. 1, 167–172
- [2] Blair, D. E., *Geometry of manifolds with structural group  $\mathcal{U}(n) \times \mathcal{O}(s)$* , J. Differential Geom., **4** (1970), 155–167
- [3] Blair, D. E., *Riemannian Geometry of Contact and Symplectic Manifolds*, 2nd. Ed., Progress in Mathematics, Vol. 209, Birkhäuser, New York, 2010
- [4] Blair, D. E and Oubiña, J. A., *Conformal and related changes of metric on the product of two almost contact metric manifolds*, Publ. Mat., **34** (1990), 199–207
- [5] Falcitelli, M. and Pastore, A. M.,  *$f$ -structures of Kenmotsu type*, Mediterr. J. Math., **3** (2006), 549–564
- [6] Fernández, L. M., *Variedades con  $K$ -estructuras. Subvariedades*, Ph. D. Thesis, University of Sevilla, Sevilla, 1987
- [7] Goldberg, S. I. and Yano, K., *On normal globally framed  $f$ -manifolds*, Tohoku Math. J., **22** (1970), No. 2, 362–370
- [8] Hasegawa, I., Okuyama, Y. and Abe, T., *On  $p$ -th Sasakian manifolds*, J. Hokkaido Univ. Educ. Nat. Sci. Section II A, **37** (1986), No. 1, 1–16
- [9] Olszak, Z., *Curvature properties of quasi-Sasakian manifolds*, Tensor (N.S.), **38** (1982), 19–28
- [10] O’Neill, B., *Semi-Riemannian Geometry with Applications to Relativity*, Pure and Applied Mathematics, Vol. 103, Academic Press, New York, 1983
- [11] Oubiña, J. A., *New classes of almost contact metric structures*, Publ. Mat., **32** (1985), 187–193
- [12] Suguri, S. and Nakayama, S.,  *$D$ -conformal deformations on almost contact metric structure*, Tensor (N.S.), **28** (1974), 125–129
- [13] Tanno, S., *The topology of contact Riemannian manifolds*, Illinois J. Math., **12** (1968), 700–717
- [14] Turgut Vanli, A. and Sari, R., *Generalized Kenmotsu manifolds*, Comm. Math. Appl., **7** (2016), No. 4, 311–328
- [15] Yano, K., *On a structure defined by a tensor field  $f$  of type  $(1,1)$  satisfying  $f^3 + f = 0$* , Tensor, **14**, (1963), 99–109

DEPARTAMENTO DE ECONOMÍA  
MÉTODOS CUANTITATIVOS E HISTORIA DE LA ECONOMÍA  
UNIVERSIDAD PABLO DE OLAVIDE, CTRA. DE UTRERA, KM. 1  
41013 SEVILLA, SPAIN  
E-mail address: psalerue@upo.es

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA  
FACULTAD DE MATEMÁTICAS  
UNIVERSIDAD DE SEVILLA  
C./ TARFIA, s.N. 41012 SEVILLA, SPAIN.  
E-mail address: lmfer@us.es  
E-mail address: aliciaprieto@us.es