

Dedicated to Professor Yeol Je Cho on the occasion of his retirement

Convergence of inexact orbits of monotone nonexpansive mappings

SIMEON REICH and ALEXANDER J. ZASLAVSKI

ABSTRACT. We study monotone nonexpansive self-mappings of a closed and convex cone in an ordered Banach space with particular emphasis on the asymptotic behavior of their inexact iterates.

1. INTRODUCTION AND MAIN RESULTS

During more than fifty years now, there has been a lot of activity regarding the fixed point theory of nonexpansive (that is, 1-Lipschitz) mappings. See, for example, [13, 14, 15, 20, 21] and the references cited therein. This activity stems from Banach's classical theorem [4] regarding the existence of a unique fixed point for a strict contraction. Since that seminal result, many developments have taken place in this area. We mention, for instance, existence results for fixed points of nonexpansive mappings which are not strictly contractive [13, 14, 18, 19]. Such results were obtained for general nonexpansive mappings in special Banach spaces, while for self-mappings of general complete metric spaces existence results were established for, the so-called, contractive mappings [17]. For general nonexpansive mappings in general Banach spaces the existence of a unique fixed point was established in the generic sense by using the Baire category approach [6, 7, 20, 21].

Another important topic in metric fixed point theory is the convergence of (inexact) iterates of a mapping to one of its fixed points.

For example, let (X, ρ) be a metric space and let $A : X \rightarrow X$ be nonexpansive. In [5] (see also Section 2.21 of [21]), under the assumption that for every $x \in X$, the sequence $\{A^i(x)\}_{i=1}^\infty$ converges, it was shown that for each given sequence of computational errors $\{r_i\}_{i=1}^\infty \subset (0, \infty)$ satisfying

$$\sum_{i=1}^{\infty} r_i < \infty,$$

each sequence $\{x_i\}_{i=0}^\infty \subset X$ such that

$$\rho(x_{i+1}, A(x_i)) \leq r_{i+1}, \quad i = 1, 2, \dots,$$

converges to a fixed point of A . This result has found several interesting applications. It is, for instance, an important ingredient in the superiorization methodology and in the study of perturbation resilience of algorithms. See, for example, [8, 9, 10, 11] and the references mentioned therein.

In the present paper we study the asymptotic behavior of inexact iterates of monotone nonexpansive mappings – a class of nonlinear mappings which is the subject of a rapidly growing area of research [1, 12].

Received: 01.11.2017. In revised form: 25.04.2018. Accepted: 15.07.2018

2010 *Mathematics Subject Classification*. 47H07, 47H09, 47H14.

Key words and phrases. *Banach space, convex cone, inexact orbit, monotone nonexpansive mapping.*

Corresponding author: Simeon Reich; sreich@technion.ac.il

Let $(X, \|\cdot\|)$ be a Banach space ordered by a closed and convex cone $X_+ \subset X$ satisfying

$$X_+ \cap (-X_+) = \{0\}.$$

Note that for all $x, y \in X$,

$$x \leq y \text{ if and only if } y - x \in X_+.$$

For each $x \in X$ and each $r > 0$, set

$$B(x, r) := \{y \in X : \|x - y\| \leq r\}.$$

Suppose that

$$(1.1) \quad \|x\| \leq \|y\| \text{ for all } x, y \in X_+ \text{ satisfying } x \leq y.$$

In this case the cone X_+ is called *normal*.

We assume that the normal cone X_+ has an interior point $x_* \in X_+$. Then there exists $r_* > 0$ such that

$$(1.2) \quad B(x_*, r_*) \subset X_+.$$

For each $x \in X$, set

$$(1.3) \quad \|x\|_* := \inf\{\lambda \in [0, \infty) : -\lambda x_* \leq x \leq \lambda x_*\}.$$

It is clear that for any $x \in X$, $\|x\|_*$ is well defined and finite, $\|\cdot\|_*$ is a norm on X and

$$(1.4) \quad \{x \in X : \|x\|_* \leq 1\} = \{x \in X : -x_* \leq x \leq x_*\}.$$

It is well known that the norms $\|\cdot\|$ and $\|\cdot\|_*$ are equivalent. Indeed, let $x \in X$. In view of (1.1) and (1.3),

$$-\|x\|_* x_* \leq x \leq \|x\|_* x_*$$

and

$$\|x\| \leq \|x + \|x\|_* x_*\| + \| \|x\|_* x_* \| \leq 2\|x\|_* \|x_*\| + \|x\|_* \|x_*\| \leq 3\|x\|_* \|x_*\|.$$

On the other hand, by (1.2),

$$B(0, r_*) + x_* \subset X_+,$$

$$B(0, r_*) \subset \{z \in X : -x_* \leq z \leq x_*\} = \{z \in X : \|z\|_* \leq 1\}$$

and for all $z \in X \setminus \{0\}$,

$$\|r_* \|z\|^{-1} z\|_* \leq 1, \quad \|z\|_* \leq r_*^{-1} \|z\|.$$

In the sequel we assume that

$$\|\cdot\| = \|\cdot\|_*.$$

Let a mapping $T : X_+ \rightarrow X_+$ satisfy

$$(1.5) \quad T(x) \leq T(y) \text{ for all } x, y \in X_+ \text{ satisfying } x \leq y$$

and

$$(1.6) \quad \|T(x) - T(y)\| \leq \|x - y\|$$

for all $x, y \in X_+$ satisfying $x \leq y$. Such a mapping T is said to be *monotone nonexpansive*.

In the present paper we establish the following two results.

Theorem 1.1. *Assume that for each $x \in X_+$, the sequence $\{T^i(x)\}_{i=1}^\infty$ converges and that a sequence $\{\gamma_i\}_{i=1}^\infty \subset (0, \infty)$ satisfies*

$$(1.7) \quad \sum_{i=1}^\infty \gamma_i < \infty.$$

Then each sequence $\{x_i\}_{i=0}^\infty \subset X_+$ such that

$$\|x_{i+1} - T(x_i)\| \leq \gamma_{i+1} \text{ for all integers } i \geq 0$$

also converges.

Theorem 1.2. *The mapping T is continuous.*

Note that an analog of Theorem 1.1 for nonexpansive mappings was obtained in [5]. The convergence property established in Theorem 1.1 is close (but not equivalent) to the shadowing property which is of interest in the qualitative study of dynamical systems [2, 16]. Applications of monotone nonexpansive mappings to the study of certain classes of matrix equations, differential equations and integral equations are discussed in [3].

2. PROOF OF THEOREM 1.1

Let a sequence $\{x_i\}_{i=0}^\infty \subset X_+$ satisfy

$$(2.8) \quad \|x_{i+1} - T(x_i)\| \leq \gamma_{i+1} \text{ for all integers } i \geq 0.$$

It is sufficient to show that $\{x_i\}_{i=0}^\infty$ is a Cauchy sequence.

Let $\epsilon > 0$ be given. In view of (1.7), there exists a natural number n_0 such that

$$(2.9) \quad \sum_{i=n_0}^\infty \gamma_i < \epsilon/4.$$

Set

$$(2.10) \quad y_{n_0} := x_{n_0}.$$

By (1.4) and (2.8),

$$(2.11) \quad -\gamma_{n_0+1}x_* \leq x_{n_0+1} - T(x_{n_0}) \leq \gamma_{n_0+1}x_*.$$

Set

$$(2.12) \quad y_{n_0+1} := T(x_{n_0}) + \gamma_{n_0+1}x_*.$$

It follows from (2.10) and (2.12) that

$$(2.13) \quad y_{n_0+1} = T(y_{n_0}) + \gamma_{n_0+1}x_*.$$

Equations (2.11) and (2.12) imply that

$$(2.14) \quad x_{n_0+1} \leq y_{n_0+1}.$$

By (2.11) and (2.12), we have

$$(2.15) \quad y_{n_0+1} - x_{n_0+1} \leq T(x_{n_0}) + \gamma_{n_0+1}x_* - T(x_{n_0}) + \gamma_{n_0+1}x_* \leq 2\gamma_{n_0+1}x_*.$$

For each integer $k \geq n_0 + 1$, set

$$(2.16) \quad y_{k+1} := T(y_k) + \gamma_{k+1}x_*.$$

We claim that for each integer $k \geq n_0 + 1$,

$$(2.17) \quad y_k \geq T^{k-n_0}(y_{n_0})$$

and

$$(2.18) \quad y_k - T^{k-n_0}(y_{n_0}) \leq \sum_{i=n_0+1}^k \gamma_i x_*.$$

In view of (2.13), inequalities (2.17) and (2.18) are valid for $k = n_0 + 1$. Assume now that $k \geq n_0 + 1$ is an integer and that (2.17) and (2.18) are true. By (2.16) and (2.17),

$$(2.19) \quad y_{k+1} = T(y_k) + \gamma_{k+1}x_* \geq T(T^{k-n_0}(y_{n_0})) = T^{k+1-n_0}(y_{n_0}).$$

It follows from (1.3), (1.6), (2.17) and (2.18) that

$$\|T(y_k) - T(T^{k-n_0}(y_{n_0}))\| \leq \|y_k - T^{k-n_0}(y_{n_0})\| \leq \sum_{i=n_0+1}^k \gamma_i$$

and

$$(2.20) \quad T(y_k) \leq T^{k+1-n_0}(y_{n_0}) + \sum_{i=n_0+1}^k \gamma_i x_*$$

Equations (2.16) and (2.20) imply that

$$(2.21) \quad \begin{aligned} y_{k+1} &= T(y_k) + \gamma_{k+1} x_* \leq T^{k+1-n_0}(y_{n_0}) + \sum_{i=n_0+1}^k \gamma_i x_* + \gamma_{k+1} x_* \\ &= T^{k+1-n_0}(y_{n_0}) + \sum_{i=n_0+1}^{k+1} \gamma_i x_* \end{aligned}$$

In view of (2.19) and (2.21), inequalities (2.17) and (2.18) hold for $k+1$ too. Thus we have shown by induction that (2.17) and (2.18) indeed hold for all integers $k \geq n_0 + 1$.

By (1.3), (2.9), (2.17) and (2.18), for all integers $k \geq n_0 + 1$, we have

$$(2.22) \quad \|y_k - T^{k-n_0}(y_{n_0})\| \leq \sum_{i=n_0+1}^{\infty} \gamma_i < \epsilon/4.$$

Next we claim that for each integer $k \geq n_0 + 1$, we have

$$(2.23) \quad x_k \leq y_k$$

and

$$(2.24) \quad y_k - x_k \leq 2 \sum_{i=n_0+1}^k \gamma_i x_*.$$

In view of (2.11) and (2.12), inequalities (2.23) and (2.24) do hold for $k = n_0 + 1$.

Assume that $k \geq n_0 + 1$ is an integer and that (2.23) and (2.24) hold. By (1.3) and (2.8),

$$(2.25) \quad -\gamma_{k+1} x_* \leq x_{k+1} - T(x_k) \leq \gamma_{k+1} x_*.$$

It follows from (2.16), (2.23) and (2.25) that

$$(2.26) \quad x_{k+1} \leq T(x_k) + \gamma_{k+1} x_* \leq T(y_k) + \gamma_{k+1} x_* = y_{k+1}.$$

Equations (2.16) and (2.25) imply that

$$(2.27) \quad \begin{aligned} y_{k+1} - x_{k+1} &= T(y_k) + \gamma_{k+1} x_* - x_{k+1} \\ &\leq T(y_k) + \gamma_{k+1} x_* - T(x_k) + \gamma_{k+1} x_* \end{aligned}$$

By (1.3), (1.6), (2.23) and (2.24),

$$\|T(y_k) - T(x_k)\| \leq \|y_k - x_k\| \leq 2 \sum_{i=n_0+1}^k \gamma_i$$

and

$$(2.28) \quad T(y_k) - T(x_k) \leq 2 \sum_{i=n_0+1}^k \gamma_i x_*.$$

It follows from (2.27) and (2.28) that

$$(2.29) \quad y_{k+1} - x_{k+1} \leq 2 \sum_{i=n_0+1}^k \gamma_i x_* + 2\gamma_{k+1} x_*$$

In view of (2.26) and (2.29), inequalities (2.23) and (2.24) are valid for $k + 1$ too. Thus we have shown by induction that (2.23) and (2.24) are indeed valid for all integers $k \geq n_0 + 1$.

By (1.3), (2.9), (2.23) and (2.24), for all integers $k \geq n_0 + 1$, we have

$$(2.30) \quad \|y_k - x_k\| \leq 2 \sum_{i=n_0+1}^k \gamma_i < \epsilon/2.$$

It now follows from (2.22) and (2.30) that for all integers $k \geq n_0 + 1$,

$$(2.31) \quad \|x_k - T^{k-n_0}(y_{n_0})\| < 3\epsilon/4.$$

The assumptions of the theorem imply that the limit

$$\lim_{k \rightarrow \infty} T^k(y_{n_0})$$

exists. When combined with (2.31), this implies that for all sufficiently large natural numbers k ,

$$\|x_k - \lim_{i \rightarrow \infty} T^i(y_{n_0})\| < \epsilon.$$

Since ϵ is an arbitrary positive number, we conclude that $\{x_k\}_{k=0}^\infty$ is a Cauchy sequence. Thus the sequence $\{x_k\}_{k=0}^\infty$ indeed converges, as asserted. Theorem 1.1 is proved.

3. PROOF OF THEOREM 1.2

Let $x_0 \in X_+$ and $\epsilon > 0$ be given. Assume that a point $x \in X_+$ satisfies

$$(3.32) \quad \|x - x_0\| \leq \epsilon/4.$$

By (1.3) and (3.32), we have

$$-(\epsilon/4)x_* \leq x - x_0 \leq (\epsilon/4)x_*$$

and

$$(3.33) \quad x_0 - (\epsilon/4)x_* \leq x \leq x_0 + (\epsilon/4)x_*.$$

It follows from (1.3), (1.5), (1.6), (3.32) and (3.33) that

$$T(x) \leq T(x_0 + 4^{-1}\epsilon x_*)$$

and

$$(3.34) \quad \begin{aligned} \|T(x) - T(x_0 + 4^{-1}\epsilon x_*)\| &\leq \|x_0 + 4^{-1}\epsilon x_* - x\| \\ &\leq 4^{-1}\epsilon + \|x_0 - x\| \leq 2^{-1}\epsilon. \end{aligned}$$

By (1.3) and (1.6), we have

$$(3.35) \quad \|T(x_0) - T(x_0 + 4^{-1}\epsilon x_*)\| \leq \|4^{-1}\epsilon x_*\| = 4^{-1}\epsilon.$$

In view of (3.34) and (3.35), we conclude that

$$\|T(x) - T(x_0)\| \leq \epsilon.$$

This completes the proof of Theorem 1.2.

Acknowledgments. The first author was partially supported by the Israel Science Foundation (Grants no. 389/12 and 820/17), by the Fund for the Promotion of Research at the Technion and by the Technion General Research Fund. Both authors are grateful to the referees for their useful comments.

REFERENCES

- [1] Alfuraidan, M. R. and Khamsi, M. A., *A fixed point theorem for monotone asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc., **146** (2018), 2451–2456
- [2] Aoki, N. and Hiraide, K., *Topological Theory of Dynamical Systems. Recent Advances*, North-Holland Mathematical Library, 52, North-Holland Publishing Co., Amsterdam, 1994
- [3] Bachar, M. and Khamsi, M. A., *Recent contributions to fixed point theory of monotone mappings*, J. Fixed Point Theory Appl., **19** (2017), 1953–1976
- [4] Banach, S., *Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales*, Fund. Math., **3** (1922), 133–181
- [5] Butnariu, D., Reich, S. and Zaslavski, A. J., *Convergence to fixed points of inexact orbits of Bregman-monotone and of nonexpansive operators in Banach spaces*, in *Fixed Point Theory and its Applications*, Yokohama Publishers, Yokohama, 2006, 11–32
- [6] de Blasi, F. S. and Myjak, J., *Sur la convergence des approximations successives pour les contractions non linéaires dans un espace de Banach*, C. R. Acad. Sci. Paris, **283** (1976), 185–187
- [7] de Blasi, F. S. and Myjak, J., *Sur la porosité de l'ensemble des contractions sans point fixe*, C. R. Acad. Sci. Paris, **308** (1989), 51–54
- [8] Censor, Y., Davidi, R. and Herman, G. T., *Perturbation resilience and superiorization of iterative algorithms*, Inverse Problems, **26** (2010), 12 pp.
- [9] Censor, Y., Davidi, R., Herman, G. T., Schulte, R. W. and Tetruashvili, L., *Projected subgradient minimization versus superiorization*, J. Optim. Theory Appl., **160** (2014), 730–747
- [10] Censor, Y. and Zaslavski, A. J., *Convergence and perturbation resilience of dynamic string-averaging projection methods*, Comput. Optim. Appl., **54** (2013), 65–76
- [11] Censor, Y. and Zaslavski, A. J., *Strict Fejér monotonicity by superiorization of feasibility-seeking projection methods*, J. Optim. Theory Appl., **165** (2015), 172–187
- [12] Espinola R. and Wiśnicki, A., *The Knaster-Tarski theorem versus monotone nonexpansive mappings*, preprint
- [13] Goebel, K. and Kirk, W. A., *Topics in Metric Fixed Point Theory*, Cambridge University Press, Cambridge, 1990
- [14] Goebel, K. and Reich, S., *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*, Marcel Dekker, New York and Basel, 1984
- [15] Kirk, W. A., *Contraction mappings and extensions*, in *Handbook of Metric Fixed Point Theory*, Kluwer, Dordrecht, 2001, 1–34
- [16] Pilyugin, S. Y., *Shadowing in Dynamical Systems*, Lecture Notes in Mathematics, 1706, Springer, Berlin, 1999
- [17] Rakotch, E., *A note on contractive mappings*, Proc. Amer. Math. Soc., **13** (1962), 459–465
- [18] Reich, S., *The fixed point property for nonexpansive mappings*, Amer. Math. Monthly, **83** (1976), 266–268
- [19] Reich, S., *The fixed point property for nonexpansive mappings, II*, Amer. Math. Monthly, **87** (1980), 292–294
- [20] Reich, S. and Zaslavski, A. J., *Generic aspects of metric fixed point theory*, in *Handbook of Metric Fixed Point Theory*, Kluwer, Dordrecht, 2001, 557–575
- [21] Reich, S. and Zaslavski, A. J., *Genericity in Nonlinear Analysis*, Developments in Mathematics, vol. **34**, Springer, New York, 2014

DEPARTMENT OF MATHEMATICS

THE TECHNION –ISRAEL INSTITUTE OF TECHNOLOGY

32000 HAIFA, ISRAEL

E-mail address: sreich@technion.ac.il

E-mail address: ajzasl@technion.ac.il