

Dedicated to Professor Yeol Je Cho on the occasion of his retirement

Iterative algorithms for split common fixed points of demicontractive operators without priori knowledge of operator norms

YONGHONG YAO^{1,2}, JEN-CHIH YAO³, YEONG-CHENG LIOU⁴ and MIHAI POSTOLACHE⁵

ABSTRACT. The split common fixed points problem for demicontractive operators has been studied in Hilbert spaces. An iterative algorithm is considered and the weak convergence result is given under some mild assumptions.

1. INTRODUCTION

Let \mathcal{H}_1 and \mathcal{H}_2 be two real Hilbert spaces equipped up their own inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$. Let $\mathcal{S}: \mathcal{H}_1 \rightarrow \mathcal{H}_1$ and $\mathcal{T}: \mathcal{H}_2 \rightarrow \mathcal{H}_2$ be two nonlinear operators. Denote the fixed point sets of \mathcal{S} and \mathcal{T} by $\text{Fix}(\mathcal{S})$ and $\text{Fix}(\mathcal{T})$, respectively. Let $\mathcal{A}: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ be a bounded linear operator with its adjoint \mathcal{A}^* .

The present article focusses on the split common fixed point problem which is to find a point $u^\dagger \in \mathcal{H}_1$ such that

$$(1.1) \quad u^\dagger \in \text{Fix}(\mathcal{S}) \quad \text{and} \quad \mathcal{A}u^\dagger \in \text{Fix}(\mathcal{T}).$$

The split common fixed point problem (1.1) is a generalization of the split feasibility problem arising from signal processing and image restoration ([4, 6, 17, 18, 19, 21]), which is to find a point u^\dagger such that

$$(1.2) \quad u^\dagger \in \mathcal{C} \quad \text{and} \quad \mathcal{A}u^\dagger \in \mathcal{Q},$$

where $\mathcal{C} \subset \mathcal{H}_1$ and $\mathcal{Q} \subset \mathcal{H}_2$ are two nonempty closed convex sets. Problem (1.1) was firstly introduced by Censor and Segal [7]. Note that solving (1.1) can be translated to solve the fixed point equation

$$x^* = \mathcal{S}(x^* - \tau \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x^*), \quad \tau > 0.$$

Whereafter, Censor and Segal proposed an algorithm for directed operators. Since then, there has been growing interest in the split common fixed point problem ([1, 3, 5, 8, 12, 13, 14, 20, 22]). In particular, Wang [16] introduced the following new iterative algorithm for the split common fixed point problem of firmly-nonexpansive mappings.

Algorithm 1.1. Choose an arbitrary initial guess x_0 .

Step 1. Given x_n , compute the next iteration via the formula:

$$(1.3) \quad x_{n+1} = x_n - \rho_n [x_n - \mathcal{S}x_n + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n], \quad n \geq 0.$$

Received: 17.09.2017. In revised form: 13.06.2018. Accepted: 15.07.2018

2010 *Mathematics Subject Classification.* 47J25, 47J20, 49N45, 65J15.

Key words and phrases. *split common fixed point, demicontractive operator, iterative algorithm, weak convergence.*

Corresponding author: Yeong-Cheng Liou; simplex.liou@hotmail.com

Step 2. If the following equality

$$(1.4) \quad \|x_{n+1} - \mathcal{S}x_{n+1} + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_{n+1}\| = 0.$$

holds, then stop; otherwise go to step 1.

Subsequently, Wang [16] demonstrated the convergence analysis of Algorithm 1.1.

Theorem 1.1. *Assume the following conditions are satisfied:*

- (A1) \mathcal{A} is a bounded linear operator;
- (A2) the solution set of problem (1.1), denoted by Ω , is nonempty;
- (A3) both \mathcal{S} and \mathcal{T} are firmly nonexpansive operators.

Let $\{x_n\}$ be the sequence generated by Algorithm 1.1. If the sequence $\{\rho_n\}$ satisfies $\sum_{n=0}^{\infty} \rho_n = \infty$ and $\sum_{n=0}^{\infty} \rho_n^2 < \infty$, then $\{x_n\}$ converges weakly to a solution z of problem (1.1), where $z = \lim_{n \rightarrow \infty} \text{proj}_{\Omega} x_n$.

At the same time, Wang[16] gave the following remark.

Remark 1.1. It is readily seen that, in Algorithm 1.1, the selection of the stepsize does not depend on the operator norm $\|\mathcal{A}\|$. It seems that the assumption (A3) cannot weaken to directed operators.

Inspired by the work in the literature, the main purpose of this paper is to give an answer to Remark 1.1. We will extend Wang's [16] result from the firmly nonexpansive operators to a more general demicontractive operators. We demonstrate the convergence of Algorithm 1.1 for solving the split common fixed point problem (1.1). Weak convergence theorem is given under some mild assumptions.

2. PRELIMINARIES

Let \mathcal{C} be a nonempty closed convex subset of a real Hilbert space \mathcal{H} .

Definition 2.1. An operator $\mathcal{T}: \mathcal{C} \rightarrow \mathcal{C}$ is said to be nonexpansive if

$$\|\mathcal{T}u - \mathcal{T}u^\dagger\| \leq \|u - u^\dagger\|$$

for all $u, u^\dagger \in \mathcal{C}$.

Definition 2.2. An operator $\mathcal{T}: \mathcal{C} \rightarrow \mathcal{C}$ is said to be firmly nonexpansive if

$$\|\mathcal{T}u - \mathcal{T}u^\dagger\|^2 \leq \|u - u^\dagger\|^2 - \|(I - \mathcal{T})u - (I - \mathcal{T})u^\dagger\|^2$$

for all $u, u^\dagger \in \mathcal{C}$.

A typical example of firmly nonexpansive operators is an orthogonal projection $\text{proj}_{\mathcal{C}}$ from \mathcal{H} onto \mathcal{C} defined by

$$\text{proj}_{\mathcal{C}}(u) := \arg \min_{v \in \mathcal{C}} \|u - v\|, u \in \mathcal{H}.$$

The metric projection $\text{proj}_{\mathcal{C}}$ of \mathcal{H} onto \mathcal{C} is characterized by

$$(2.5) \quad \langle u - \text{proj}_{\mathcal{C}}(u), v - \text{proj}_{\mathcal{C}}(u) \rangle \geq 0$$

for all $u \in \mathcal{H}, v \in \mathcal{C}$.

Definition 2.3. An operator $\mathcal{T}: \mathcal{C} \rightarrow \mathcal{C}$ is said to be directed if

$$\|\mathcal{T}u - u^\dagger\|^2 \leq \|u - u^\dagger\|^2 - \|\mathcal{T}u - u\|^2$$

for all $u \in \mathcal{C}$ and $u^\dagger \in \text{Fix}(\mathcal{T})$.

The class of directed operators is an important class since it includes the orthogonal projections and the subgradient projectors which are fundamental in the convex optimization.

Definition 2.4. An operator \mathcal{T} is called demicontractive if there exists a constant $\beta \in [0, 1)$ such that

$$\|\mathcal{T}u - v\|^2 \leq \|u - v\|^2 + \beta\|u - \mathcal{T}u\|^2,$$

or equivalently,

$$(2.6) \quad \langle u - \mathcal{T}u, u - v \rangle \geq \frac{1 - \beta}{2} \|u - \mathcal{T}u\|^2,$$

for all $(u, v) \in \mathcal{H} \times \text{Fix}(\mathcal{T})$.

Remark 2.2. It is clear that the demicontractive operators include the directed operators as special cases. The class of demicontractive operators is fundamental because many common types of operators arising in optimization belong to this class, see for example [?] and references therein.

Definition 2.5. An operator \mathcal{T} is said to be demiclosed if for any sequence $\{x_n\}$ which weakly converges to \tilde{x} , and if the sequence $\{\mathcal{T}(x_n)\}$ strongly converges to z , then $\mathcal{T}(\tilde{x}) = z$.

Definition 2.6. A sequence $\{x_n\}$ is called Fejér-monotone with respect to a given nonempty set Ω if for every $x \in \Omega$,

$$\|x_{n+1} - x\| \leq \|x_n - x\|, \quad \text{for all } n \geq 0.$$

Next we adopt the following notations:

- $x_n \rightharpoonup x$ means that x_n converges weakly to x ;
- $\omega_w(x_n) := \{x : \exists x_{n_j} \rightharpoonup x\}$ is the weak ω -limit set of the sequence $\{x_n\}$.

Lemma 2.1. ([2]) Let Ω be a nonempty closed convex subset in \mathcal{H} . If the sequence $\{x_n\}$ is Fejér monotone with respect to Ω , then we have the following conclusions:

- (i) $x_n \rightharpoonup x^\dagger \in \Omega$ iff $\omega_w(x_n) \subset \Omega$;
- (ii) the sequence $\{\text{proj}_\Omega(x_n)\}$ converges strongly;
- (iii) if $x_n \rightharpoonup x^\dagger \in \Omega$, then $x^\dagger = \lim_{n \rightarrow \infty} \text{proj}_\Omega(x_n)$.

Lemma 2.2. ([15]) Let $\{a_n\}$ and $\{b_n\}$ be positive real sequences such that $\sum_{n=0}^\infty b_n < \infty$. If either $a_{n+1} \leq (1 + b_n)a_n$ or $a_{n+1} \leq a_n + b_n$, then $\lim_{n \rightarrow \infty} a_n$ exists.

3. MAIN RESULTS

In this section, we will give the convergence analysis of Algorithm 1.1 for solving the split common fixed points problem (1.1). For the purpose, the following hypotheses are involved.

- (HP1): \mathcal{H}_1 and \mathcal{H}_2 are two real Hilbert spaces;
- (HP2): $\mathcal{S}: \mathcal{H}_1 \rightarrow \mathcal{H}_1$ and $\mathcal{T}: \mathcal{H}_2 \rightarrow \mathcal{H}_2$ are two demicontractive operators with coefficients $\beta \in [0, 1)$ and $\mu \in [0, 1)$, respectively;
- (HP3): $\mathcal{S}: \mathcal{H}_1 \rightarrow \mathcal{H}_1$ and $\mathcal{T}: \mathcal{H}_2 \rightarrow \mathcal{H}_2$ are Lipschitz continuous with Lipschitz constant $L > 1$;
- (HP4): $\mathcal{A}: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is a bounded linear operator with its adjoint operator \mathcal{A}^* .

We use Ω to denote the solution set of problem (1.1), that is,

$$\Omega = \{\bar{z} : \bar{z} \in \text{Fix}(\mathcal{S}) \text{ and } \mathcal{A}\bar{z} \in \text{Fix}(\mathcal{T})\}.$$

Throughout, assume Ω is nonempty.

First, we have the following remark.

Remark 3.3. If (1.4) holds, then x_{n+1} solves problem (1.1). As a matter of fact, we have the following a more general conclusion.

Proposition 3.1. z^\dagger solves (1.1) iff $\|z^\dagger - \mathcal{S}z^\dagger + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z^\dagger\| = 0$.

Proof. If z^\dagger solves (1.1), then $z^\dagger = \mathcal{S}z^\dagger$ and $(I - \mathcal{T})\mathcal{A}z^\dagger = 0$. It is obvious that $\|z^\dagger - \mathcal{S}z^\dagger + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z^\dagger\| = 0$.

To see the converse, assume that $\|z^\dagger - \mathcal{S}z^\dagger + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z^\dagger\| = 0$, then for any $z \in \Omega$, we obtain

$$\begin{aligned}
 (3.7) \quad 0 &= \|z^\dagger - \mathcal{S}z^\dagger + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z^\dagger\| \|z^\dagger - z\| \\
 &\geq \langle z^\dagger - \mathcal{S}z^\dagger + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z^\dagger, z^\dagger - z \rangle \\
 &= \langle z^\dagger - \mathcal{S}z^\dagger, z^\dagger - z \rangle + \langle \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z^\dagger, z^\dagger - z \rangle \\
 &= \langle z^\dagger - \mathcal{S}z^\dagger, z^\dagger - z \rangle + \langle (I - \mathcal{T})\mathcal{A}z^\dagger, \mathcal{A}z^\dagger - \mathcal{A}z \rangle.
 \end{aligned}$$

Since \mathcal{S} and \mathcal{T} are demicontractive, from (2.6), we deduce

$$(3.8) \quad \langle z^\dagger - \mathcal{S}z^\dagger, z^\dagger - z \rangle \geq \frac{1 - \beta}{2} \|z^\dagger - \mathcal{S}z^\dagger\|^2,$$

and

$$(3.9) \quad \langle (I - \mathcal{T})\mathcal{A}z^\dagger, \mathcal{A}z^\dagger - \mathcal{A}z \rangle \geq \frac{1 - \mu}{2} \|(I - \mathcal{T})\mathcal{A}z^\dagger\|^2.$$

By (3.7)-(3.9), we get

$$\begin{aligned}
 (3.10) \quad 0 &\geq \langle z^\dagger - \mathcal{S}z^\dagger + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z^\dagger, z^\dagger - z \rangle \\
 &\geq \frac{1 - \beta}{2} \|z^\dagger - \mathcal{S}z^\dagger\|^2 + \frac{1 - \mu}{2} \|(I - \mathcal{T})\mathcal{A}z^\dagger\|^2.
 \end{aligned}$$

Since $\beta, \mu \in [0, 1)$, we deduce $z^\dagger \in \text{Fix}(\mathcal{S})$ and $\mathcal{A}z^\dagger \in \text{Fix}(\mathcal{T})$ by (3.10). Therefore, z^\dagger solves problem (1.1). The proof is completed. \square

We assume that the sequence $\{x_n\}$ generated by Algorithm 1.1 is infinite. In other words, Algorithm 1.1 does not terminate in a finite number of iterations. In this case, we demonstrate the convergence of the sequence $\{x_n\}$ generated by Algorithm 1.1.

Theorem 3.2. Assume the following conditions are satisfied

- (i) $I - \mathcal{S}$ and $I - \mathcal{T}$ are demiclosed at zero;
- (ii) $\sum_{n=0}^\infty \rho_n = \infty$ and $\sum_{n=0}^\infty \rho_n^2 < \infty$.

Then the sequence $\{x_n\}$ generated by Algorithm 1.1 converges weakly to a solution z^* of problem (1.1), where $z^* = \lim_{n \rightarrow \infty} \text{proj}_\Omega(x_n)$.

Proof. Firstly, we prove that the sequence $\{x_n\}$ is bounded. Let $z \in \Omega$. Set $y_n = x_n - \mathcal{S}x_n + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n$ for all $n \geq 0$. From (2.6), we have

$$\begin{aligned}
 \langle y_n, x_n - z \rangle &= \langle x_n - \mathcal{S}x_n + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n, x_n - z \rangle \\
 &= \langle x_n - \mathcal{S}x_n, x_n - z \rangle + \langle \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n, x_n - z \rangle \\
 &\geq \frac{1 - \beta}{2} \|x_n - \mathcal{S}x_n\|^2 + \frac{1 - \mu}{2} \|(I - \mathcal{T})\mathcal{A}x_n\|^2 \\
 &= \frac{1 - \beta}{2} \|x_n - \mathcal{S}x_n\|^2 + \frac{(1 - \mu)\|\mathcal{A}\|^2}{2\|\mathcal{A}\|^2} \|(I - \mathcal{T})\mathcal{A}x_n\|^2 \\
 &\geq \frac{1 - \beta}{2} \|x_n - \mathcal{S}x_n\|^2 + \frac{(1 - \mu)}{2\|\mathcal{A}\|^2} \|A^*(I - \mathcal{T})\mathcal{A}x_n\|^2 \\
 (3.11) \quad &\geq \frac{\min\{1 - \beta, 1 - \mu\}}{2 \max\{1, \|\mathcal{A}\|^2\}} (\|x_n - \mathcal{S}x_n\|^2 + \|A^*(I - \mathcal{T})\mathcal{A}x_n\|^2) \\
 &\geq \frac{\min\{1 - \beta, 1 - \mu\}}{4 \max\{1, \|\mathcal{A}\|^2\}} (\|x_n - \mathcal{S}x_n\| + \|A^*(I - \mathcal{T})\mathcal{A}x_n\|)^2 \\
 &\geq \frac{\min\{1 - \beta, 1 - \mu\}}{4 \max\{1, \|\mathcal{A}\|^2\}} \|x_n - \mathcal{S}x_n + A^*(I - \mathcal{T})\mathcal{A}x_n\|^2 \\
 &= \frac{\min\{1 - \beta, 1 - \mu\}}{4 \max\{1, \|\mathcal{A}\|^2\}} \|y_n\|^2 \\
 &= \tau \|y_n\|^2,
 \end{aligned}$$

where $\tau = \frac{\min\{1 - \beta, 1 - \mu\}}{4 \max\{1, \|\mathcal{A}\|^2\}}$.

According to (1.3) and (3.11), we derive

$$\begin{aligned}
 (3.12) \quad \|x_{n+1} - z\|^2 &= \|x_n - z + \rho_n y_n\|^2 \\
 &= \|x_n - z\|^2 - 2\rho_n \langle y_n, x_n - z \rangle + \rho_n^2 \|y_n\|^2 \\
 &\leq \|x_n - z\|^2 - 2\rho_n \tau \|y_n\|^2 + \rho_n^2 \|y_n\|^2.
 \end{aligned}$$

Since S and T are L -Lipschitzian, we have

$$\begin{aligned}
 (3.13) \quad \|y_n\| &= \|x_n - \mathcal{S}x_n + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n - [z - \mathcal{S}z + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z]\| \\
 &\leq \|(I - \mathcal{S})x_n - (I - \mathcal{S})z\| + \|\mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n - \mathcal{A}^*(I - \mathcal{T})\mathcal{A}z\| \\
 &\leq (L + 1)\|x_n - z\| + (L + 1)\|\mathcal{A}\|^2 \|x_n - z\| \\
 &= (L + 1)(1 + \|\mathcal{A}\|^2) \|x_n - z\|.
 \end{aligned}$$

By (3.12) and (3.13), we get

$$\begin{aligned}
 (3.14) \quad \|x_{n+1} - z\|^2 &\leq \|x_n - z\|^2 - 2\rho_n \tau \|y_n\|^2 + \rho_n^2 \|y_n\|^2 \\
 &\leq \|x_n - z\|^2 + \rho_n^2 (L + 1)^2 (1 + \|\mathcal{A}\|^2)^2 \|x_n - z\|^2 - 2\rho_n \tau \|y_n\|^2 \\
 &\leq \|x_n - z\|^2 + \rho_n^2 (L + 1)^2 (1 + \|\mathcal{A}\|^2)^2 \|x_n - z\|^2.
 \end{aligned}$$

Noting that $\sum_{n=0}^\infty \rho_n^2 < \infty$, from Lemma 2.2 and (3.14), we deduce $\lim_{n \rightarrow \infty} \|x_n - z\|$ exists. Hence, $\{x_n\}$ is bounded and so is $\{y_n\}$.

Next, we show that every weak cluster point of the sequence $\{x_n\}$ belongs to the solution set of problem (1.1), i.e., $\omega_w(x_n) \subset \Omega$.

From (3.12), we obtain

$$\begin{aligned}
 2\tau \rho_n \|y_n\|^2 &\leq \|x_n - z\|^2 - \|x_{n+1} - z\|^2 + \rho_n^2 \|y_n\|^2 \\
 &\leq \|x_n - z\|^2 - \|x_{n+1} - z\|^2 + \rho_n^2 M,
 \end{aligned}$$

where $M = \sup_n \{\|y_n\|^2\}$.

An induction induces that

$$(3.15) \quad 2\tau \sum_{n=0}^{\infty} \rho_n \|y_n\|^2 \leq \|x_0 - z\|^2 + M \sum_{n=0}^{\infty} \rho_n^2 < \infty,$$

which implies that

$$(3.16) \quad \liminf_{n \rightarrow \infty} \|y_n\| = 0$$

due to the fact that $\sum_{n=0}^{\infty} \rho_n = \infty$.

Observe that

$$(3.17) \quad \begin{aligned} \|y_{n+1} - y_n\| &= \|x_{n+1} - \mathcal{S}x_{n+1} + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_{n+1} \\ &\quad - [x_n - \mathcal{S}x_n + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n]\| \\ &\leq \|x_{n+1} - \mathcal{S}x_{n+1} - [x_n - \mathcal{S}x_n]\| \\ &\quad + \|\mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_{n+1} - \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n\| \\ &\leq (L + 1)\|x_{n+1} - x_n\| + (L + 1)\|\mathcal{A}\|^2\|x_{n+1} - x_n\| \\ &= (L + 1)(1 + \|\mathcal{A}\|^2)\|x_{n+1} - x_n\| \\ &= (L + 1)(1 + \|\mathcal{A}\|^2)\rho_n\|y_n\|. \end{aligned}$$

It follows that

$$(3.18) \quad \begin{aligned} \|y_{n+1}\|^2 &= \|y_n\|^2 + 2\langle y_n, y_{n+1} - y_n \rangle + \|y_{n+1} - y_n\|^2 \\ &\leq \|y_n\|^2 + 2\|y_n\|\|y_{n+1} - y_n\| + \|y_{n+1} - y_n\|^2 \\ &\leq \|y_n\|^2 + 2(L + 1)(1 + \|\mathcal{A}\|^2)\rho_n\|y_n\|^2 + (L + 1)^2(1 + \|\mathcal{A}\|^2)^2\rho_n^2\|y_n\|^2. \end{aligned}$$

From (3.15), we know that $\sum_{n=0}^{\infty} \rho_n \|y_n\|^2 < \infty$. At the same time, $\sum_{n=0}^{\infty} \rho_n^2 < \infty$ by the assumption. Thus, we can apply Lemma 2.2 to (3.18) to deduce $\lim_{n \rightarrow \infty} \|y_n\|$ exists. This together with (3.16) implies that

$$(3.19) \quad \lim_{n \rightarrow \infty} \|y_n\| = \lim_{n \rightarrow \infty} \|x_n - \mathcal{S}x_n + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n\| = 0.$$

Observe that

$$(3.20) \quad \begin{aligned} &\frac{1 - \beta}{2}\|x_n - \mathcal{S}x_n\|^2 + \frac{1 - \mu}{2}\|(I - \mathcal{T})\mathcal{A}x_n\|^2 \\ &\leq \langle x_n - \mathcal{S}x_n, x_n - z \rangle + \langle \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n, x_n - z \rangle \\ &= \langle x_n - \mathcal{S}x_n + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n, x_n - z \rangle \\ &\leq \|x_n - \mathcal{S}x_n + \mathcal{A}^*(I - \mathcal{T})\mathcal{A}x_n\|\|x_n - z\|. \end{aligned}$$

In view of (3.19) and (3.20), we have

$$(3.21) \quad \lim_{n \rightarrow \infty} \|x_n - \mathcal{S}x_n\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|(I - \mathcal{T})\mathcal{A}x_n\| = 0.$$

By the demiclosedness (at zero) of $I - \mathcal{S}$ and $I - \mathcal{T}$, we deduce immediately $\omega_w(x_n) \subset \Omega$. To this end, the conditions of Lemma 2.1 are all satisfied. Consequently, $x_n \rightarrow z^* = \lim_{n \rightarrow \infty} \text{proj}_{\Omega}(x_n)$. The proof is completed. \square

Since the demicontractive operators include the directed operators and the firmly-nonexpansive mappings operators as special cases, we have immediately the following corollaries.

Corollary 3.1. *Assume the following conditions are satisfied*

- (i) \mathcal{H}_1 and \mathcal{H}_2 are two real Hilbert spaces;
- (ii) $\mathcal{S}: \mathcal{H}_1 \rightarrow \mathcal{H}_1$ and $\mathcal{T}: \mathcal{H}_2 \rightarrow \mathcal{H}_2$ are two directed operators;

- (iii) $S : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ and $\mathcal{T} : \mathcal{H}_2 \rightarrow \mathcal{H}_2$ are Lipschitz continuous with Lipschitz constant $L > 1$;
- (iv) $I - S$ and $I - \mathcal{T}$ are demiclosed at zero;
- (v) $\sum_{n=0}^{\infty} \rho = \infty$ and $\sum_{n=0}^{\infty} \rho_n^2 < \infty$.

Then the sequence $\{x_n\}$ generated by Algorithm 1.1 converges weakly to a solution z^* of problem (1.1), where $z^* = \lim_{n \rightarrow \infty} \text{proj}_{\Omega}(x_n)$.

Corollary 3.2. Assume the following conditions are satisfied

- (i) \mathcal{H}_1 and \mathcal{H}_2 are two real Hilbert spaces;
- (ii) $S : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ and $\mathcal{T} : \mathcal{H}_2 \rightarrow \mathcal{H}_2$ are two firmly-nonexpansive mappings;
- (iii) $\sum_{n=0}^{\infty} \rho_n = \infty$ and $\sum_{n=0}^{\infty} \rho_n^2 < \infty$.

Then the sequence $\{x_n\}$ generated by Algorithm 1.1 converges weakly to a solution z^* of problem (1.1), where $z^* = \lim_{n \rightarrow \infty} \text{proj}_{\Omega}(x_n)$.

Acknowledgements.

Yonghong Yao was supported by the grant TD13-5033. Yeong-Cheng Liou was supported in part by the grants KMU-Q107011 and MOST 106-2218-E-040-001-MY2. Jen-Chih Yao was partially supported by the Grant MOST 106-2923-E-039-001-MY3.

REFERENCES

- [1] Ansari, Q. H., Rehan, A. and Wen C. F., *Implicit and explicit algorithms for split common fixed point problems*, J. Nonlinear Convex Anal., **17** (2016), 1381–1397
- [2] Bauschke, H. H. and Borwein, J. M., *On projection algorithms for solving convex feasibility problems*, SIAM Review., **38** (1996), 367–426
- [3] Boikanyo, O. A., *A strongly convergent algorithm for the split common fixed point problem*, Appl. Math. Comput., **265** (2015), 844–853
- [4] Byrne, C., *A unified treatment of some iterative algorithms in signal processing and image reconstruction*, Inverse Probl., **20** (2004), 103–120
- [5] Cegielski, A., *General method for solving the split common fixed point problem*, J. Optim. Theory Appl., **165** (2015), 385–404
- [6] Ceng, L. C., Ansari Q. H. and Yao, J. C., *An extragradient method for split feasibility and fixed point problems*, Comput. Math. Appl., **64** (2012), 633–642
- [7] Censor, Y. and Segal, A., *The split common fixed point problem for directed operators*, J. Convex Anal., **16** (2009), 587–600
- [8] Kraikaew P. and Saejung, S., *On split common fixed point problems*, J. Math. Anal. Appl., **415** (2014), 513–524
- [9] Mărușter, S. and Rus, I. A., *Kannan contractions and strongly demicontractive mappings*, Creat. Math. Inform., **24** (2015), No. 2, 171–180
- [10] Mărușter, S., *A note on the convergence of Mann iteration*, Creat. Math. Inform., **26** (2017), No. 1, 85–88
- [11] Maingé, P.-E. and Mărușter, S., *Convergence in norm of modified Krasnoselski-Mann iterations for fixed points of demicontractive mappings*, Appl. Math. Comput., **217** (2011), No. 24, 9864–9874
- [12] Moudafi, A., *The split common fixed-point problem for demicontractive mappings*, Inverse Probl., **26** (2010), Art. ID 055007
- [13] Takahashi, W., *The split common fixed point problem and strong convergence theorems by hybrid methods in two Banach spaces*, J. Nonlinear Convex Anal., **17** (2016), 1051–1067
- [14] Takahashi, S. and Takahashi, W., *The split common null point problem and the shrinking projection method in two Banach spaces*, Linear Nonlinear Anal., **1** (2015), 297–304
- [15] Tan, K. K. and Xu, H. K., *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, J. Math. Anal. Appl., **178** (1993), 301–308
- [16] Wang, F., *A new method for split common fixed-point problem without priori knowledge of operator norms*, J. Fixed Point Theory Appl., DOI 10.1007/s11784-017-0434-0
- [17] Wang, F. and Xu, H. K., *Cyclic algorithms for split feasibility problems in Hilbert spaces*, Nonlinear Anal., **74** (2011), 4105–4111
- [18] Xu, H. K., *Iterative methods for the split feasibility problem in infinite-dimensional Hilbert spaces*, Inverse Probl., **26** (2010), Art. ID 105018
- [19] Yao, Y. H., Leng, L. M., Postolache, M. and Zheng, X. X., *A unified framework for the two-sets split common fixed point problem in Hilbert spaces*, J. Nonlinear Sci. Appl., **9** (2016), 6113–6125

- [20] Yao, Y. H., Liou, Y. C. and Yao, J. C., *Split common fixed point problem for two quasi-pseudo-contractive operators and its algorithm construction*, Fixed Point Theory Appl., **2015** (2015), Art. ID 127
- [21] Yao, Y. H., Liou, Y. C. and Yao, J. C., *Iterative algorithms for the split variational inequality and fixed point problems under nonlinear transformations*, J. Nonlinear Sci. Appl., **10** (2017), 843–854
- [22] Zheng, X. X., Yao, Y. H., Liou, Y. C. and Leng, L. M., *Fixed point algorithms for the split problem of demicontractive operators*, J. Nonlinear Sc. Appl., **10** (2017), 1263–1269

^{1,2}NORTH MINZU UNIVERSITY
SCHOOL OF MATHEMATICS AND INFORMATION SCIENCE
YINCHUAN, 750021 CHINA

DEPARTMENT OF MATHEMATICS
TIANJIN POLYTECHNIC UNIVERSITY
TIANJIN 300387 CHINA
E-mail address: yaoyonghong@aliyun.com

³ CHINA MEDICAL UNIVERSITY
CENTER FOR GENERAL EDUCATION
TAICHUNG, 40402 TAIWAN
E-mail address: yaojc@mail.cmu.edu.tw

⁴ KAOHSIUNG MEDICAL UNIVERSITY
DEPARTMENT OF HEALTHCARE ADMINISTRATION AND MEDICAL INFORMATICS
CENTER FOR BIG DATA ANALYTICS & INTELLIGENT HEALTHCARE
RESEARCH CENTER OF NONLINEAR ANALYSIS AND OPTIMIZATION
DEPARTMENT OF MEDICAL RESEARCH
KAOHSIUNG, 807 TAIWAN
E-mail address: simplex.liou@hotmail.com

⁵ DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE
UNIVERSITY "POLITEHNICA" OF BUCHAREST
SPLAIUL INDEPENDENTEI 313, 060042 BUCHAREST, ROMANIA
E-mail address: emscolar@yahoo.com