

In memoriam Prof. Charles E. Chidume (1947-2021)

Inertial Iteration Scheme for Approximating Fixed Points of Lipschitz Pseudocontractive Maps in Arbitrary Real Banach Spaces

P.U. NWOKORO, D. F. AGBEBAKU, E.E. CHIMA, A.C. ONAH, O.U. OGUGUO,
AND M.O. OSILIKE

ABSTRACT. We study a perturbed inertial Krasnoselskii-Mann-type algorithm and prove that the algorithm is an approximate fixed point sequence for Lipschitz pseudocontractive maps in arbitrary real Banach spaces. Strong convergence results are then established for our inertial iteration scheme for approximation of fixed points of Lipschitz pseudocontractive maps and solutions of certain important accretive-type operator equations in certain real Banach spaces. Implementation of our algorithm is illustrated using numerical examples in both finite and infinite dimensional Banach spaces. Our results improve rate of convergence and extend several related recent results.

1. INTRODUCTION

Let X be a real Banach with dual X^* . Let $J : X \rightarrow X^*$ denote the normalized duality mapping given by

$$Jx := \{f^* \in X^* : \langle x, f^* \rangle = \|x\|^2 = \|f^*\|^2\},$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing (see for example [10]). J is single-valued if X^* is strictly convex (see for example [10]) and in what follows we denote single-valued normalized duality mapping by j . A mapping $T : D(T) \subseteq X \rightarrow R(T) \subseteq X$ is said to be L -Lipschitzian (see for example [10]) if there exists $L \geq 0$ such that

$$\|Tx - Ty\| \leq L\|x - y\|, \quad \forall x, y \in D(T). \quad (1.1)$$

T is said to be a *contraction* if $L \in [0, 1)$ and T is said to be *nonexpansive* if $L = 1$ (see for example [3, 6, 10, 28]). T is said to be *pseudocontractive* (see for example [10]) if

$$\|x - y\| \leq \|x - y + \lambda((x - Tx) - (y - Ty))\|, \quad \forall x, y \in D(T), \lambda > 0. \quad (1.2)$$

An operator $A : D(A) \subseteq X \rightarrow R(A) \subseteq X$ is said to be *accretive* (see for example [10]) if for all $t > 0$ and every $x, y \in D(A)$ we have

$$\|x - y\| \leq \|x - y + t(Ax - Ay)\|. \quad (1.3)$$

It follows from Lemma 1.1 of Kato [15] that T is pseudocontractive if and only if for all $x, y \in D(T)$ there exists $j(x - y) \in J(x - y)$ such

$$\langle (I - T)x - (I - T)y, j(x - y) \rangle \geq 0. \quad (1.4)$$

Received: 24.07.2021. In revised form: 10.12.2021. Accepted: 14.12.2021

2010 *Mathematics Subject Classification.* 47H09, 47H25, 47J25, 65J15.

Key words and phrases. Lipschitz pseudocontractive mappings, Inertial terms, Weak Convergence, Strong convergence, Accretive operators.

Corresponding author: M.O. Osilike; micah.osilike@unn.edu.ng

Furthermore, A is accretive if and only if for each $x, y \in D(A)$, there exists $j(x - y) \in J(x - y)$ such that

$$\langle Ax - Ay, j(x - y) \rangle \geq 0. \quad (1.5)$$

An accretive operator A is said to be m -accretive (see for example [10]) if $R(I + \lambda A) = X$, $\forall \lambda > 0$, where $R(I + \lambda A)$ is the range of $(I + \lambda A)$. The class of nonexpansive maps is a proper subclass of the class of pseudocontractive maps and T is pseudocontractive if and only if $(I - T)$ is accretive. Other important subclasses of pseudocontractive maps are the class of *strictly pseudocontractive operators* and the class of *strongly pseudocontractive operators*. T is strictly pseudocontractive if $\forall x, y \in D(T)$ and for some $k \in [0, 1)$, we have

$$\langle (I - T)x - (I - T)y, j(x - y) \rangle \geq k\|(I - T)x - (I - T)y\|^2, \quad (1.6)$$

and T is strongly pseudocontractive if

$$\langle (I - T)x - (I - T)y, j(x - y) \rangle \geq k\|x - y\|^2, \quad \forall x, y \in D(T) \text{ and for some } k \in [0, 1), \quad (1.7)$$

(see for example [10]). The classes of strictly pseudocontractions and strongly pseudocontractions are independent and every strictly pseudocontraction is Lipschitz (see for example [3, 10]).

In the iterative approximation of fixed points of nonexpansive maps, strictly pseudocontractive maps and strongly pseudocontractive maps, the iterative scheme of Mann [22]:

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T x_n, \quad n \geq 1; \quad (1.8)$$

where $\{\alpha_n\}_{n=1}^{\infty}$ is a suitable sequence in $[0, 1]$ have played pivotal role. In [8] Chidume and Mutangadura showed with an example in the 2-dimensional Euclidean plane that the original Mann iteration may fail to converge to a fixed point of a Lipschitz pseudocontractive selfmap T defined on a nonempty compact subset. For the class of pseudocontractive maps T , the Ishikawa iteration [14]:

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T[(1 - \beta_n)x_n + \beta_n T x_n], \quad n \geq 1, \quad (1.9)$$

and its modifications have played very important role in the iterative approximation of fixed points of T when $F(T) \neq \emptyset$. However, the Ishikawa iteration scheme for Lipschitz pseudocontractions yield only weak convergence usually obtained mostly from $\lim_{n \rightarrow \infty} \|x_n - T x_n\| = 0$; and require "compactness" assumption either on the operator or the domain of the operator or even both to yield strong convergence. Most of the related results are also confined to Hilbert spaces. Often, very strong conditions are imposed on the fixed-point set, $F(T)$ to obtain strong convergence using the usual Ishikawa iteration process (see for example [3, 10, 30, 36]). For instance in [30], the author required that $F(T)$ is finite where T is a continuous pseudocontractive-type self-mapping of a nonempty convex compact of a Hilbert space, and in [36] the authors required that the interior of $F(T)$ is nonempty where T is a Lipschitz pseudocontractive self-mapping of a nonempty closed convex subset of a Hilbert space. Thus many other schemes have been recently studied by several authors to achieve relatively fast strong convergence with mild assumptions on the operator, its domain, its set of fixed points and other necessary components (see for example [1, 2, 3, 7, 10, 11, 12, 13, 16, 17, 18, 19, 20, 21, 23, 26, 27, 29, 31, 34, 35]). The Krasnoselskii-Mann and certain modifications and other prominent algorithms like the Halpern-type algorithms had been used to obtain weak and strong convergence results. In [9] the authors studied the following algorithm in arbitrary real Banach spaces:

Algorithm CZ ([9]). Let X be a real Banach space, K a nonempty closed convex subset of X and $T : K \rightarrow K$ a given operator. Then for arbitrary $x_1 \in K$ the sequence $\{x_n\}$ is generated by

$$x_{n+1} = (1 - \lambda_n)x_n + \lambda_n T x_n - \lambda_n \theta_n (x_n - x_1), \quad n \geq 1, \quad (1.10)$$

where $\{\lambda_n\}$ and $\{\theta_n\}$ are real sequences in $(0, 1)$ such that: (i) $\lambda_n(1 + \theta_n) < 1$, (ii) $\lim_{n \rightarrow \infty} \theta_n = 0$, (iii) $\sum_{n=1}^{\infty} \lambda_n \theta_n = \infty$, (iv) $\lim_{n \rightarrow \infty} \frac{\lambda_n}{\theta_n} = 0$, (v) $\lim_{n \rightarrow \infty} \frac{\theta_{n-1} - \theta_n}{\lambda_n \theta_n^2} = 0$. They proved that the sequence $\{x_n\}$ is an approximate fixed point sequence for Lipschitz pseudocontractive maps, T , in arbitrary real Banach spaces (i.e., $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$). They further obtained strong convergence of the sequence to $p \in F(T)$ when $F(T) \neq \emptyset$ in certain real Banach spaces much more general than Hilbert spaces. The algorithm has been studied extensively in recent years by various authors; it appears simpler than the Ishikawa iteration scheme and the results are proved in general Banach spaces. However, the implementation of the algorithm shows it as being very slow and hence there is a need to accelerate the scheme to achieve relatively fast convergence rate.

It is our purpose in this paper to introduce appropriate inertial term in Algorithm CZ to accelerate the convergence. We consider the following:

Algorithm 1.1. For arbitrary $x_0, x_1 \in X$, the sequence $\{x_n\}$ is generated by

$$\begin{cases} w_n = x_n + \theta_n(x_n - x_{n-1}), & n \geq 1 \\ x_{n+1} = (1 - \lambda_n)w_n + \lambda_n T w_n - \lambda_n \sigma_n(w_n - x_0), & n \geq 1, \end{cases} \tag{1.11}$$

where $\{\lambda_n\}$ and $\{\sigma_n\}$ are suitable sequences in $(0, 1)$; for some suitable $\theta \in (0, 1)$, $0 \leq \theta_n \leq \bar{\theta}_n$, where

$$\bar{\theta}_n = \begin{cases} \min\{\theta, \frac{\epsilon_n}{\|x_n - x_{n-1}\|}\}, & x_n \neq x_{n-1} \\ \theta, & \text{otherwise,} \end{cases}$$

and $\{\epsilon_n\} \subseteq \mathbb{R}^+$ is a suitable sequence such that $\sum_{n=1}^{\infty} \epsilon_n < \infty$. We prove that under suitable conditions on $\{\lambda_n\}, \{\sigma_n\}, \{\theta_n\}, \{\epsilon_n\}$ and appropriate θ , the perturbed inertial Mann algorithm (1.11) is an approximate fixed point sequence for Lipschitz pseudocontractive maps in arbitrary real Banach spaces.

Interesting fast strong convergence results which generalize and improve several related recent results are proved. Implementation of the algorithm is demonstrated in finite and infinite dimensional Banach spaces with comparison with Algorithm CZ, the scheme without inertial term.

2. PRELIMINARIES

We shall need the following results:

Lemma 2.1. ([15]) *Let X be an arbitrary Banach space, X^* the dual of X and let $x, y \in X$ be arbitrary. Then $\|x\| \leq \|x + \lambda y\|, \forall \lambda > 0$ if and only if there exists $j(x) \in J(x)$ such that $\operatorname{Re}\langle y, j(x) \rangle \geq 0$.*

Lemma 2.2. ([24]) *Let $\{a_n\}_{n=1}^{\infty}, \{\alpha_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$ be sequences of nonnegative real numbers and let $\psi : [0, \infty) \rightarrow [0, \infty)$ be a strictly increasing continuous function with $\psi(0) = 0, \psi(t) > 0, t \in (0, \infty)$. Let*

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$,
 - (ii) $\sum_{n=1}^{\infty} \alpha_n = \infty$,
 - (iii) $\lim_{n \rightarrow \infty} \frac{b_n}{\alpha_n} = 0$,
 - (iv) $a_{n+1}^2 \leq a_n^2 - \alpha_n \psi(a_{n+1}) + b_n$.
- Then $\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 2.3. ([25]) *Let X be a Banach space, K a nonempty closed convex subset of X and $T : K \rightarrow X$ a continuous pseudocontractive mapping satisfying the weakly inward condition. Then for each $x_0 \in K$, there exists a unique path $t \rightarrow y_t \in K; t \in [0, 1)$, satisfying*

$$y_t = tT y_t + (1 - t)x_0. \tag{2.1}$$

Remark 2.1. If in Lemma 2.3 $F(T) \neq \emptyset$, then $\{y_t\}$ is bounded. Furthermore, if either (i) X has a uniformly Gâteaux differentiable norm and is such that every closed convex bounded subset of K has the fixed point property (FPP) for nonexpansive self-mappings, or (ii) X is a reflexive and strictly convex Banach space with a uniformly Gâteaux differentiable norm, then as $t \rightarrow 1$, the path converges strongly to a fixed point of T (see [25, 9]).

3. MAIN RESULTS

We now prove the following results:

Theorem 3.1. Let X be an arbitrary real Banach space and let $T : X \rightarrow X$ be an L -Lipschitzian pseudocontractive mapping with $F(T) \neq \emptyset$. Let $\{\lambda_n\}$ and $\{\sigma_n\}$ be real sequence in $(0, 1)$ such that (i) $\lambda_n(1 + \sigma_n) < 1$, (ii) $\lim_{n \rightarrow \infty} \sigma_n = 0$, (iii) $\sum_{n=1}^{\infty} \lambda_n \sigma_n = \infty$, (iv) $\sum_{n=1}^{\infty} \lambda_n^2 < \infty$; (v) $\lim_{n \rightarrow \infty} \frac{|\sigma_n - \sigma_{n-1}|}{\lambda_n \sigma_n^2} = 0$, (vi) $\lim_{n \rightarrow \infty} \frac{\lambda_n}{\sigma_n} = 0$. Let $\{\epsilon_n\} \subseteq \mathbb{R}^+$ be such that $\sum_{n=1}^{\infty} \epsilon_n < \infty$, $\lim_{n \rightarrow \infty} \frac{\epsilon_n}{\lambda_n \sigma_n} = 0$, and let $\theta \in [0, 1)$. For arbitrary $x_0, x_1 \in X$, let $\{x_n\}$ be the sequence generated by (1.11):

$$\begin{cases} w_n = x_n + \theta_n(x_n - x_{n-1}), & n \geq 1 \\ x_{n+1} = (1 - \lambda_n)w_n + \lambda_n T w_n - \lambda_n \sigma_n(w_n - x_0), & n \geq 1, \end{cases}$$

where $\theta \in [0, 1)$, $0 \leq \theta_n \leq \bar{\theta}_n$ and

$$\bar{\theta}_n = \begin{cases} \min\{\theta, \frac{\epsilon_n}{\|x_n - x_{n-1}\|}\}, & x_n \neq x_{n-1} \\ \theta, & \text{otherwise.} \end{cases} \tag{3.1}$$

Then $\lim_{n \rightarrow \infty} \|x_n - T x_n\| = 0$.

Proof. We divide the proof as follows:

Claim 1: The sequences $\{x_n\}_{n=1}^{\infty}$, $\{w_n\}_{n=1}^{\infty}$, $\{T x_n\}_{n=1}^{\infty}$ and $\{T w_n\}_{n=1}^{\infty}$ are bounded.

Let $p \in F(T)$ be arbitrary and define $G_n : X \rightarrow X$ by

$G_n x := T x - \sigma_n x + \sigma_n x_0$, $n \geq 1$. Then $G_n p = (1 - \sigma_n)p + \sigma_n x_0$ and

$$\langle (I - G_n)x - (I - G_n)y, j(x - y) \rangle \geq \sigma_n \|x - y\|^2, \forall x, y \in X, n \geq 1. \tag{3.2}$$

From (3.2) we obtain

$$\langle (I - G_n - \sigma_n I)x - (I - G_n - \sigma_n I)y, j(x - y) \rangle \geq 0, \forall x, y \in X, n \geq 1, \tag{3.3}$$

and it follows from Lemma 2.1 that $\forall x, y \in X, n \geq 1$ we have

$$\|x - y\| \leq \|x - y + \lambda[(I - G_n - \sigma_n I)x - (I - G_n - \sigma_n I)y]\|. \tag{3.4}$$

From (1.11) we obtain

$$\begin{aligned} w_n &= (1 + \lambda_n + \lambda_n \sigma_n)x_{n+1} + \lambda_n(I - G_n - \sigma_n I)x_{n+1} - \lambda_n w_n \\ &\quad + \lambda_n(G_n x_{n+1} - G_n w_n) + 2\lambda_n^2(w_n - G_n w_n) \end{aligned} \tag{3.5}$$

Furthermore,

$$p = (1 + \lambda_n + \lambda_n \sigma_n)p + \lambda_n(I - G_n - \sigma_n I)p - \lambda_n p - \lambda_n \sigma_n(p - x_0). \tag{3.6}$$

Thus from (3.5) and (3.6) we obtain

$$\begin{aligned} \|w_n - p\| &= \|(1 + \lambda_n + \lambda_n \sigma_n)(x_{n+1} - p) + \lambda_n[(I - G_n - \sigma_n I)x_{n+1} \\ &\quad - (I - G_n - \sigma_n I)p] - \lambda_n(w_n - p) + \lambda_n(G_n x_{n+1} - G_n w_n) \\ &\quad + 2\lambda_n^2(w_n - G_n w_n) + \lambda_n \sigma_n(p - x_0)\| \\ &\geq (1 + \lambda_n + \lambda_n \sigma_n)\|x_{n+1} - p\| - \lambda_n\|w_n - p\| - \lambda_n\|G_n x_{n+1} - G_n w_n\| \\ &\quad - 2\lambda_n^2\|w_n - G_n w_n\| - \lambda_n \sigma_n\|p - x_0\|. \end{aligned}$$

Hence

$$\begin{aligned}
 \|x_{n+1} - p\| &\leq \frac{(1 + \lambda_n)}{(1 + \lambda_n + \lambda_n \sigma_n)} \|w_n - p\| + \frac{1}{(1 + \lambda_n + \lambda_n \sigma_n)} [\lambda_n \|G_n x_{n+1} - G_n w_n\| \\
 &\quad + 2\lambda_n^2 \|w_n - G_n w_n\| + \lambda_n \sigma_n \|p - x_0\|] \\
 &\leq (1 + \lambda_n)[1 - \lambda_n(1 + \sigma_n) + \lambda_n^2(1 + \sigma_n)^2] \|x_n - p\| \\
 &\quad + (1 + \lambda_n)\theta_n \|x_n - x_{n-1}\| \\
 &\quad + \frac{1}{(1 + \lambda_n + \lambda_n \sigma_n)} [\lambda_n^2(2 + L + \sigma_n) \|w_n - G_n w_n\| \\
 &\quad + \lambda_n \sigma_n \|p - x_0\|] \\
 &\leq (1 + \lambda_n)[1 - \lambda_n(1 + \sigma_n) + \lambda_n^2(1 + \sigma_n)^2] \|x_n - p\| \\
 &\quad + (1 + \lambda_n)\theta_n \|x_n - x_{n-1}\| \\
 &\quad + \frac{1}{(1 + \lambda_n + \lambda_n \sigma_n)} [\lambda_n^2(2 + L + \sigma_n)(1 + L + \sigma_n)(\|x_n - p\| \\
 &\quad + \theta_n \|x_n - x_{n-1}\|) + \sigma_n \|p - x_0\|] + \lambda_n \sigma_n \|p - x_0\| \\
 &\leq \left[1 - \lambda_n \sigma_n + \lambda_n^2[(1 + \sigma_n)(\sigma_n + \lambda_n(1 + \sigma_n)) \right. \\
 &\quad \left. + (2 + L + \sigma_n)(1 + L + \sigma_n)] \right] \|x_n - p\| \\
 &\quad + [1 + \lambda_n + \lambda_n^2(2 + L + \sigma_n)] \theta_n \|x_n - x_{n-1}\| \\
 &\quad + \lambda_n^2 \sigma_n (2 + L + \sigma_n) \|p - x_0\| + \lambda_n \sigma_n \|p - x_0\| \\
 &= [1 - \lambda_n \sigma_n + a_n] \|x_n - p\| + b_n + \lambda_n \sigma_n \|p - x_0\|, \tag{3.7}
 \end{aligned}$$

where

$$a_n = \lambda_n^2[(1 + \sigma_n)(\sigma_n + \lambda_n(1 + \sigma_n)) + (2 + L + \sigma_n)(1 + L + \sigma_n)],$$

and

$$b_n = [1 + \lambda_n + \lambda_n^2(2 + L + \sigma_n)] \theta_n \|x_n - x_{n-1}\| + \lambda_n^2 \sigma_n (2 + L + \sigma_n) \|p - x_0\|.$$

From (3.7) we obtain

$$\begin{aligned}
 \|x_{n+1} - p\| &\leq [1 - \lambda_n \sigma_n + a_n] \max\{\|x_n - p\|, \|p - x_0\|\} + b_n \\
 &\quad + \lambda_n \sigma_n \max\{\|x_n - p\|, \|p - x_0\|\} \\
 &\leq [1 + a_n] \max\{\|x_n - p\|, \|p - x_0\|\} + b_n \\
 &\quad \vdots \\
 &\leq \prod_{j=1}^n (1 + a_j) \|p - x_0\| + \sum_{j=1}^n b_j < \infty.
 \end{aligned}$$

Thus $\{x_n\}_{n=1}^\infty$, $\{w_n\}_{n=1}^\infty$, $\{Tx_n\}_{n=1}^\infty$ and $\{Tw_n\}_{n=1}^\infty$ are bounded.

Claim 2: The sequences $\{y_n\}_{n=1}^\infty$ and $\{Ty_n\}_{n=1}^\infty$, where $t_n := \frac{1}{1 + \sigma_n}$, and $y_n \equiv y_{t_n} = t_n T y_n + (1 - t_n)x_0$ are bounded.

Observe that $\lim_{n \rightarrow \infty} t_n = 1$ and since $F(T) \neq \emptyset$, it follows from Proposition 2 of [25] that $\{y_n\}$ is bounded. Since T is Lipschitz pseudocontractive, we have also that $\{Ty_n\}$ is bounded.

Claim 3: $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$.

$$\begin{aligned}
\|x_{n+1} - y_n\|^2 &\leq \|x_n - y_n + \theta_n(x_n - x_{n-1}) - \lambda_n(w_n - Tw_n) - \lambda_n\sigma_n(w_n - x_0)\|^2 \\
&\leq \|x_n - y_n\|^2 + 2\langle \theta_n(x_n - x_{n-1}) - \lambda_n(w_n - Tw_n) \\
&\quad - \lambda_n\sigma_n(w_n - x_0), j(x_{n+1} - y_n) \rangle \\
&\leq \|x_n - y_n\|^2 - 2\langle \lambda_n(w_n - Tw_n) + \lambda_n\sigma_n(w_n - x_0), j(x_{n+1} - y_n) \rangle \\
&\quad + 2\theta_n\|x_n - x_{n-1}\|\|x_{n+1} - y_n\| \\
&= \|x_n - y_n\|^2 - 2\lambda_n\sigma_n\|x_{n+1} - y_n\|^2 \\
&\quad + 2\lambda_n\langle \sigma_n(x_{n+1} - y_n) - (w_n - Tw_n) - \sigma_n(w_n - x_0), j(x_{n+1} - y_n) \rangle \\
&\quad + 2\theta_n\|x_n - x_{n-1}\|\|x_{n+1} - y_n\| \\
&= \|x_n - y_n\|^2 - 2\lambda_n\sigma_n\|x_{n+1} - y_n\|^2 \\
&\quad + 2\lambda_n\langle \sigma_n(x_{n+1} - w_n) + \sigma_n(x_0 - y_n) - (w_n - Tw_n), j(x_{n+1} - y_n) \rangle \\
&\quad + 2\theta_n\|x_n - x_{n-1}\|\|x_{n+1} - y_n\| \\
&= \|x_n - y_n\|^2 - 2\lambda_n\sigma_n\|x_{n+1} - y_n\|^2 \\
&\quad + 2\lambda_n\langle \sigma_n(x_{n+1} - w_n) + \sigma_n(x_0 - y_n) - (y_n - Ty_n) \\
&\quad - (x_{n+1} - Tx_{n+1} - (y_n - Ty_n)) + x_{n+1} - Tx_{n+1} \\
&\quad - (w_n - Tw_n), j(x_{n+1} - y_n) \rangle + 2\theta_n\|x_n - x_{n-1}\|\|x_{n+1} - y_n\| \quad (3.8)
\end{aligned}$$

Observe that $\sigma_n(x_0 - y_n) - (y_n - Ty_n) = 0$, and the pseudocontractive property of T implies that $\langle x_{n+1} - Tx_{n+1} - (y_n - Ty_n), j(x_{n+1} - y_n) \rangle \geq 0$. Thus it follows from (3.8) that

$$\begin{aligned}
\|x_{n+1} - y_n\|^2 &\leq \|x_n - y_n\|^2 - 2\lambda_n\sigma_n\|x_{n+1} - y_n\|^2 \\
&\quad + 2\lambda_n\sigma_n\|x_{n+1} - w_n\|\|x_{n+1} - y_n\| \\
&\quad + 2\lambda_n\|x_{n+1} - w_n - (Tx_{n+1} - Tw_n)\|\|x_{n+1} - y_n\| \\
&\quad + 2\theta_n\|x_n - x_{n-1}\|\|x_{n+1} - y_n\| \\
&\leq \|x_n - y_n\|^2 - 2\lambda_n\sigma_n\|x_{n+1} - y_n\|^2 \\
&\quad + [2\lambda_n\sigma_n + 2\lambda_n(1 + L)]\|x_{n+1} - w_n\|\|x_{n+1} - y_n\| \\
&\quad + 2\theta_n\|x_n - x_{n-1}\|\|x_{n+1} - y_n\| \\
&\leq \|x_n - y_{n-1}\|^2 + 2\|y_{n-1} - y_n\|\|x_n - y_n\| - 2\lambda_n\sigma_n\|x_{n+1} - y_n\|^2 \\
&\quad + \lambda_n[2\lambda_n(\sigma_n + 1 + L)\|w_n - G_n w_n\| \\
&\quad + 2\frac{\theta_n}{\lambda_n}\|x_n - x_{n-1}\|\|x_{n+1} - y_n\| \quad (3.9)
\end{aligned}$$

Since T is pseudocontractive, then $\langle (I - T)x - (I - T)y, j(x - y) \rangle \geq 0$ and hence

$$\|x - y\| \leq \|x - y + \lambda((I - T)x - (I - T)y)\|, \forall x, y \in X, \lambda > 0.$$

Thus

$$\begin{aligned}
 \|y_n - y_{n-1}\| &\leq \|y_n - y_{n-1} + \frac{1}{\sigma_n}[(I - T)y_n - (I - T)y_{n-1}]\| \\
 &= \frac{1}{\sigma_n} \|\sigma_n(y_n - y_{n-1}) + y_n - Ty_n - y_{n-1} + Ty_{n-1}\| \\
 &= \frac{1}{\sigma_n} \|(1 + \sigma_n)y_n - (1 + \sigma_n)y_{n-1} - Ty_n + Ty_{n-1}\| \\
 &= \frac{1}{\sigma_n} \|Ty_n + \frac{(1 - t_n)}{t_n}x_0 - (1 + \sigma_{n-1})y_{n-1} \\
 &\quad + (1 + \sigma_{n-1})y_{n-1} - (1 + \sigma_n)y_{n-1} - Ty_n + Ty_{n-1}\| \\
 &= \frac{1}{\sigma_n} \|\sigma_n x_0 - \sigma_{n-1}x_0 - (\sigma_n - \sigma_{n-1})y_{n-1}\| \\
 &= \frac{1}{\sigma_n} |\sigma_n - \sigma_{n-1}| \|x_0 - y_{n-1}\| \\
 &\leq \frac{1}{\sigma_n} |\sigma_n - \sigma_{n-1}| [\|x_0\| + \|y_{n-1}\|].
 \end{aligned} \tag{3.10}$$

Using (3.10) in (3.9) yields

$$\begin{aligned}
 \|x_{n+1} - y_n\|^2 &\leq \|x_n - y_{n-1}\|^2 - 2\lambda_n \sigma_n \|x_{n+1} - y_n\|^2 \\
 &\quad + 2\lambda_n \sigma_n \left[\frac{1}{\lambda_n \sigma_n^2} |\sigma_n - \sigma_{n-1}| (\|x_0\| + \|y_{n-1}\|) \|x_n - y_n\| \right. \\
 &\quad + \left[\frac{\lambda_n}{\sigma_n} (\sigma_n + 1 + L) \|w_n - G_n w_n\| \right. \\
 &\quad \left. \left. + \frac{\theta_n}{\lambda_n \sigma_n} \|x_n - x_{n-1}\| \|x_{n+1} - y_n\| \right] \right]
 \end{aligned} \tag{3.11}$$

Since $\{x_n\}, \{y_n\}, \{w_n\}, \{Tw_n\}, \{G_n w_n\}$ and $\{\sigma_n\}$ are bounded, it follows from the conditions on $\{\lambda_n\}, \{\sigma_n\}, \{\theta_n\}$ and $\{\epsilon_n\}$ and Lemma 2.2 that $\lim_{n \rightarrow \infty} \|x_n - y_{n-1}\| = 0$. Thus

$\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$. Finally we proof

Claim 4: $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$.

Since $0 \leq \|y_n - Ty_n\| = (1 - t_n)\|x_0 - Ty_n\| \rightarrow 0$ as $n \rightarrow \infty$, we obtain that

$$0 \leq \|x_n - Tx_n\| \leq (1 + L)\|x_n - y_n\| + \|y_n - Ty_n\| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

□

Prototype for our iteration parameters are: $\lambda_n = \frac{1}{(n+1)^a}; \sigma_n = \frac{1}{(n+1)^b};$
 where $0 < b < a < 1; a + b < 1; 2a > 1$ (in particular $\lambda_n = \frac{1}{2(n+1)^{\frac{3}{5}}}, \sigma_n = \frac{1}{(n+1)^{\frac{1}{5}}};$
 $\epsilon_n = \frac{1}{(n+1)^2}$, and $\theta = \frac{1}{2}$.

Theorem 3.2. *Let X be a real Banach space with a uniformly Gâteaux differentiable norm, and let $T : X \rightarrow X$ be an L -Lipschitzian pseudocontractive map with $F(T) \neq \emptyset$. Let every nonempty closed convex and bounded subset C of X have the fixed point property (FPP) for nonexpansive selfmaps and let $\{\lambda_n\}, \{\sigma_n\}, \{\epsilon_n\}, \theta, \{\theta_n\}$, and $\{\bar{\theta}_n\}$ be as in Theorem 3.1. Then the sequence $\{x_n\}$ generated from arbitrary $x_0, x_1 \in X$ by (1.11) converges strongly to a fixed point of T .*

Proof. It follows from Theorem 1 of [25] that $\{y_n\}$ converges to a fixed point $p \in F(T)$ and since $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$, we obtain that $\{x_n\}$ converges strongly to $p \in F(T)$. □

Corollary 3.1. Let X be a real uniformly smooth Banach space and let $T : X \rightarrow X$ be an L -Lipschitzian pseudocontractive map with $F(T) \neq \emptyset$. Let $\{\lambda_n\}$, $\{\sigma_n\}$, $\{\epsilon_n\}$, θ , $\{\theta_n\}$, and $\{\bar{\theta}_n\}$ be as in Theorem 3.1. Then the sequence $\{x_n\}$ generated from arbitrary $x_0, x_1 \in X$ by (1.11) converges strongly to a fixed point of T .

Proof. Every uniformly smooth Banach space has a Gâteaux differentiable norm and every nonempty closed convex bounded subset C of X has the fixed point property for nonexpansive selfmap (see for example [33]). \square

Theorem 3.3. Let X be a real reflexive and strictly convex Banach space with a uniformly Gâteaux differentiable norm, and let $T : X \rightarrow X$ be a nonexpansive map with $F(T) \neq \emptyset$. Let $\{\lambda_n\}$, $\{\sigma_n\}$, $\{\epsilon_n\}$, θ , $\{\theta_n\}$, and $\{\bar{\theta}_n\}$ be as in Theorem 3.1. Then the sequence $\{x_n\}$ generated from arbitrary $x_0, x_1 \in X$ by (1.11) converges strongly to a fixed point of T .

Proof. Follows as in the proof Theorem 3.2 using Theorem 2 of [25]. \square

Corollary 3.2. Let X be a real Banach space with a uniformly Gâteaux differentiable norm and such that every nonempty closed convex and bounded subset C of X have the fixed point property (FPP) for nonexpansive selfmaps. Let $A : X \rightarrow X$ be an accretive operator with $A^{-1}0 = N(A) = \{x \in X : Ax = 0\} \neq \emptyset$ and let $\{\lambda_n\}$, $\{\sigma_n\}$, $\{\epsilon_n\}$, θ , $\{\theta_n\}$, and $\{\bar{\theta}_n\}$ be as in Theorem 3.1. For arbitrary $x_0, x_1 \in X$, let $\{x_n\}$ be the sequence generated by

$$\begin{cases} w_n = x_n + \theta_n(x_n - x_{n-1}), & n \geq 1 \\ x_{n+1} = w_n - \lambda_n A w_n - \lambda_n \sigma_n(w_n - x_0), & n \geq 1. \end{cases} \quad (3.12)$$

Then $\{x_n\}$ converges strongly to a point $x^* \in A^{-1}0$.

Corollary 3.3. Let X be a real uniformly smooth Banach space Let $A : X \rightarrow X$ be an accretive operator with $A^{-1}0 = N(A) = \{x \in X : Ax = 0\} \neq \emptyset$ and let $\{\lambda_n\}$, $\{\sigma_n\}$, $\{\epsilon_n\}$, θ , $\{\theta_n\}$, and $\{\bar{\theta}_n\}$ be as in Theorem 3.1. Then the sequence $\{x_n\}$ generated from arbitrary $x_0, x_1 \in X$ by (3.12) converges strongly to a point $x^* \in A^{-1}0$.

Corollary 3.4. Let X be a real reflexive and strictly convex Banach space with a uniformly Gâteaux differentiable norm. Let $A : X \rightarrow X$ be an accretive operator with $A^{-1}0 = N(A) = \{x \in X : Ax = 0\} \neq \emptyset$. and let $\{\lambda_n\}$, $\{\sigma_n\}$, $\{\epsilon_n\}$, θ , $\{\theta_n\}$, and $\{\bar{\theta}_n\}$ be as in Theorem 3.1. Then the sequence $\{x_n\}$ generated from arbitrary $x_0, x_1 \in X$ by (3.12) converges strongly to a point $x^* \in A^{-1}0$.

4. NUMERICAL EXAMPLES

In this section, numerical illustration of the convergence of the iterative scheme 1.11 and iterative scheme CZ discussed in this paper are presented. The setting for the numerical example is that of a real Hilbert space. Using different examples, we show graphically and with a table of numerical values the convergence results discussed in this paper.

All codes are written in MATLAB, and implemented using an HP Elitebook 6930p computer with Pentium(R) DUAL-CORE CPU T4200 with 2.00Hz and 2GB RAM.

Example 4.1. Let \mathbb{R}^2 denote the 2-dimensional Euclidean plane. Define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$Tx = T((x_1, x_2)) = (x_1, x_2) + (x_2, -x_1) = (x_1 + x_2, x_2 - x_1),$$

for each $x = (x_1, x_2) \in \mathbb{R}^2$. Then T is 2-Lipschitz pseudocontractive.

Let $x_0 = (-0.01, -0.01)$; $x_1 = (1, 1)$; $\lambda_n = \frac{1}{(n+3)^{0.501}}$, $\sigma_n = \frac{1}{(n+3)^{0.2}}$; $\epsilon_n = \frac{1}{(n+3)^{1.01}}$, and $\theta = 0.5$. Then algorithm (1.11) converges strongly to $p = (0, 0)$ which is the only fixed point of T . It is shown in [8] that the original Mann algorithm (1.8) [22] fails to converge to the fixed point of T . Observe that algorithm (1.11) converges much faster than Algorithm CZ (see Table 1 and Figure 1 below).




FIGURE 1. Graph showing the convergence of iterative schemes

No. of iter.		Iterative scheme 1.11		Iterative scheme CZ	
n	Time (Secs.)	Sequence $x_n = (x_n^1, x_n^2)$		Sequence $x_n = (x_n^1, x_n^2)$	
			$\ x_{n+1} - x_n\ $		$\ x_{n+1} - x_n\ $
1	0.181851	(1.214830, 0.129164)	0.896943	(1.499307, 0.500693)	0.706127
2	0.202965	(0.849417, -0.533430)	0.756676	(1.561283, -0.007158)	0.511619
592	1.300140	(-0.003305, 0.001887)	0.000001	(0.330551, -0.189437)	0.000124
593	1.304350	(-0.003304, 0.001886)	0.000001	(0.330432, -0.189404)	0.000124
2537	4.359200	(-0.002412, 0.001589)	0.000000	(0.241209, -0.159220)	0.000022
2538	4.361230	(-0.002412, 0.001589)	0.000000	(0.241188, -0.159211)	0.000022
4992	8.265880	(-0.002082, 0.001446)	0.000000	(0.208156, -0.144830)	0.000010
4993	8.267110	(-0.002082, 0.001446)	0.000000	(0.208147, -0.144826)	0.000010

TABLE 1. Table showing some terms of the sequence generated by iterative scheme 1.11 and CZ, values of $\|x_{n+1} - x_n\|$ and CPU time for the indicated values of n

Example 4.2. Let $X = \ell_2(\mathbb{R}) = \{x = \{x_i\}_{i=1}^\infty : x_i \in \mathbb{R} \text{ and } \sum_{i=1}^\infty |x_i|^2 < \infty\}$. Define $T : X \rightarrow X$ by $Tx = (0, -3x_2, -3x_3, \dots)$. Then T is $\frac{1}{2}$ - strictly pseudocontractive and hence Lipschitz pseudocontractive. Let $x_0 = (-0.01, -0.01, -0.01, 0, 0, 0, \dots)$; $x_1 = (1, 1, 1, 0, 0, 0, \dots)$, $\lambda_n = \frac{1}{(n+3)^{0.51}}$, $\sigma_n = \frac{1}{(n+3)^{0.2}}$; $\epsilon_n = \frac{1}{(n+3)^{1.01}}$, and $\theta = 0.1$. Then algorithm (3.1) converges strongly to $p = (0, 0, 0, \dots)$ which is the only fixed point of T . Observe that algorithm (1.11) converges much faster than Algorithm CZ (see Table 2 and Figure 2 below).

In the two examples given above the stopping criteria is $\|x_{n+1} - x_n\| \leq 10^{-7}$. This implies that the error of approximating the fixed point of the given maps is negligible. From Figure 1 and Table 1, it is clearly seen that as consecutive terms of the sequence get close enough (as close as a difference of 10^{-7}), the sequence generated is seen to approach the fixed point of the map which is 0. A lower stopping will make no much




FIGURE 2. Graph showing the convergence of iterative schemes

No. of iter.		Iterative scheme 1.11	Iterative scheme CZ
n	Time (Secs.)	$\ x_{n+1} - x_n\ $	$\ x_{n+1} - x_n\ $
1	0.006275	3.618540	2.832740
2	0.008063	4.572730	3.311260
548	0.679860	0.000001	0.000071
549	0.680848	0.000001	0.000071
1999	2.489850	0.000000	0.000016
2000	2.490580	0.000000	0.000016
3083	3.872630	0.000000	0.000010
3084	3.873390	0.000000	0.000010

TABLE 2. Table showing some terms of the sequence generated by iterative scheme 1.11 and CZ, values of $\|x_{n+1} - x_n\|$ and CPU time for the indicated values of n

difference while a higher stopping criteria will truncate the computation too early which might leads to higher computational error. Clearly, the error = $\|x_n - 0\|$ tends to 0 as n tends to ∞ which shows that the convergence of the sequence generated by the algorithms converges. The same holds for Figure 2 and Table 2.

ACKNOWLEDGEMENT

P.U. Nwokoro, D.F. Agbebaku, E.E. Chima, A.C. Onah and Osuo-siseken Uzoma Ogugua received assistance from the facilities of Pastor E.A. Adeboye Professorial Chair. They are very thankful to the Chair.

REFERENCES

- [1] Alvarez, F. and Attouch, H., *An inertial proximal method for maximal monotone operators via discretization of a nonlinear oscillator with damping*, Set-valued Anal. **9** (2001), 3-11.
- [2] Beck, A. and Teboulle, M., *A fast iterative shrinkage-thresholding algorithm for linear inverse problems*, SIAM J. Imaging Sci. **2** (1) (2009), 183-202.
- [3] Berinde, V., *Iterative Approximation of Fixed Points*, Lectures Notes 1912 Springer 2002.
- [4] Bertsekas, D. P. and Tsitsiklis, J. N., *Parallel and Distributed Computations: Numerical Methods* Athena Scientific Belmont, MA, USA, 1997.
- [5] Browder, F. E. and Petryshyn, W. V., *Construction of fixed points of nonlinear mappings in Hilbert spaces*, J. Math. Anal. Appl. **20** (1967), 197-228.
- [6] Browder, F. E., *Nonexpansive Nonlinear Operators in Banach Spaces*, Proc. Nat. Acad. Sci, **54** (1965), 1041-1044.
- [7] Chambolle, A. and Dossal, C. H., *On the convergence of the iterates of the "fast iterative shrinkage/thresholding algorithm*, J. Optim. Theory Appl. **166** (2015), 968-982.
- [8] Chidume, C. E. and Mutangadura, S. A., *An example on the Mann iteration method for Lipschitz pseudocontractions* Proc. Amer. Math. Soc. **129** (8), (2001) 2359-2363.
- [9] Chidume, C. E. and Zegeye, H., *Approximate fixed point sequences and convergence theorems for Lipschitz pseudocontractive maps*, Proc. Amer. Math. Soc. **132** (3) (2003), 831-840.
- [10] Chidume, C. E., *Geometric Properties of Banach Spaces and Nonlinear Iterations*, Lecture Notes in Mathematics 1965, Springer 2009.
- [11] Dong, Q. L., Jiang, D., Cholamjiak, P. and Shehu, Y., *A strong convergence result involving an inertial forward-backward algorithm for monotone inclusions*, J. Fixed Point Theory Appl. **19** (2017), 3097-3118.
- [12] Dong, Q. L., Yuan, H. B., Cho, Y. J. and Rassias, T. M., *Modified inertial Mann algorithm and inertial CQ-algorithm for nonexpansive mappings*, Optim. Lett. **12** (1)(2018) 87-102.
- [13] Hussain, N., Ullah, K. and Arshad, M., *Fixed point approximation of Suzuki generalized nonexpansive mapping via new faster iteration process*, arXiv:1802.09888v1[math.FA] 2018.
- [14] Ishikawa, S., *Fixed point by a new iteration method*, Proc. Amer. Math. Soc. **44** (1) (1974), 147-150.
- [15] Kato, T., *Nonlinear semigroups and evolution equations*, J. Math. Soc. Japan **19** (1967), 508-520.
- [16] Kim, T. H. and Xu, H. K., *Strong convergence of modified Mann iterations*, Nonlinear Anal., **61** (2005) 51-60.
- [17] Li, M. and Yao, Y., *Strong convergence of an Iterative algorithm for λ -strictly pseudocontractive mappings in Hilbert spaces*, An. Șt. Univ. Ovidius Constanța **18**(1), (2010), 219-228.
- [18] Mainge, P. E., *Strong convergence of projected subgradient methods for nonsmooth and nonstrictly convex minimization*, Set-Valued Analysis, **16** (7) (2008) 899-912.
- [19] Lorenz, D. A., *Constructing test instances for basis pursuit denoising*, IEEE Trans. Signal Process. **61** (2013), 1210-1214.
- [20] Mainge, P. E. and Maruster, S., *Convergence in Norm of modified Krasnoselkii-Mann iterations for fixed points of demicontractive mappings*, Appl. Maths. and Comp., **217** (2011), 9864-9874.
- [21] Majee, P. and Nahak, C., *A modified iterative method for a finite collection of non-self mappings and family of variational inequality problems*, Med. J. Math **15:58** (2018).
- [22] Mann, W. R., *Mean, value methods in iteration*, Proceedings of the American Mathematical Society, **4** (3), (1953), 506-510.
- [23] Marino, G. and Xu, H. K., *Weak and strong convergence theorems for strict pseudocontractions in Hilbert spaces*, J. Math. Anal. Appl., **329** (2007), 336-349.
- [24] Moore, C. and Nnoli, B. V. C., *Iterative solution of nonlinear equations involving set-valued uniformly accretive operators*, Computers and Mathematics with applications **42** (2001), 131-140.
- [25] Morales, C. H. and Jung, J. S., *Convergence of paths for pseudo-contractive mappings in Banach spaces*, Pro. Amer. Math. Soc. **128** (11) (2000), 3411-3419.
- [26] Moudafi, A. and Oliny, M., *Convergence of a splitting inertial proximal method for monotone operators*, J. Comput. Appl. Math. **155** (2003), 447-454.
- [27] Nakajo, K. and Takahashi, W., *Strong convergence theorems for nonexpansive mappings and nonexpansive semigroups*, J. Math. Anal. Appl. **279** (2003), 372-379.
- [28] Opial, Z., *Weak convergence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591-597.
- [29] Polyak, B. T., *Some methods of speeding up the convergence of iteration methods*, Zh. Vychisl. Mat. Mat. Fiz. **4** (1964), 1-17.
- [30] Qihou, L., *The convergence theorems of the sequence of Ishikawa iterates for hemicontractive mappings*, J. Math. Anal. Appl. **148** (1990), 55-62.
- [31] Shehu, Y., Iyiola, O. S. and Ogbuisi, F. U., *Iterative method with inertial terms for nonexpansive mappings: applications to compressed sensing*, Numerical Algorithms **83** (2020) 1321-1347.
- [32] Shehu, Y. and Gilbali, A., *Inertial Krasnoselkii-Mann method in Banach spaces*, Mathematics (2020) **8** 638.

- [33] Turett, B., *A dual view of a theorem of Baillon*, *Nonlinear Analysis and Applications* (S.P. Singh and J.H. Burry, eds), Marcel Dekker, New York, 1982, pp. 279-286.
- [34] Ullah, K. and Arshad, M., *New three step iteration process and fixed point approximation in Banach spaces*, *J. Linear & Topological Algebra* **7** (2) (2018), 87-100.
- [35] Yao, Y., Zhou H. and Liou, Y-C., *Strong convergence of a modified Krasnoselski-Mann iterative algorithm for non-expansive mappings*, *J. Appl. Math. Comput.* **29**, (2009), 383-389.
- [36] Zegeye, H., Shahzad, N. and Alghamdi, M. A., *Convergence of Ishikawa's iteration method for pseudocontractive mappings*, *Nonlinear Analysis* **74** (2011) 7304-7311.

UNIVERSITY OF NIGERIA, NSUKKA
DEPARTMENT OF MATHEMATICS
4100010, NIGERIA
Email address: peter.nwokoro@unn.edu.ng

UNIVERSITY OF NIGERIA, NSUKKA
DEPARTMENT OF MATHEMATICS
4100010, NIGERIA
Email address: dennis.agbebaku@unn.edu.ng

BINGHAM UNIVERSITY, KARU
DEPARTMENT OF MATHEMATICS
NIGERIA
Current address: Fixed Point Theory and Application Research Group (FPTA-RG), Department of Mathematics, University of Nigeria, Nsukka, 4100010, Nigeria
Email address: epukeezinneuk@yahoo.com; chima.eucharia@binghamuni.edu.ng

UNIVERSITY OF NIGERIA, NSUKKA
DEPARTMENT OF MATHEMATICS
4100010, NIGERIA
Email address: onah87@gmail.com

UNIVERSITY OF NIGERIA, NSUKKA
DEPARTMENT OF MATHEMATICS
4100010, NIGERIA
Email address: dokufunmi@yahoo.com

UNIVERSITY OF NIGERIA, NSUKKA
DEPARTMENT OF MATHEMATICS
4100010, NIGERIA
Email address: micah.osilike@unn.edu.ng