

In memoriam Prof. Charles E. Chidume (1947-2021)

A hybrid scheme for fixed points of a countable family of generalized nonexpansive-type maps and finite families of variational inequality and equilibrium problems, with applications

MARKJOE O. UBA, MARIA A. ONYIDO, CYRIL I. UDEANI, AND PETER U. NWOKORO

ABSTRACT. Let C be a nonempty closed and convex subset of a uniformly smooth and uniformly convex real Banach space E with dual space E^* . We present a novel hybrid method for finding a common solution of a family of equilibrium problems, a common solution of a family of variational inequality problems and a common element of fixed points of a family of a general class of nonlinear nonexpansive maps. The sequence of this new method is proved to converge strongly to a common element of the families. Our theorem and its applications complement, generalize, and extend various results in literature.

1. INTRODUCTION

Let E be a real Banach space with topological dual E^* . Let $C \subset E$ be closed and convex with J_C also closed and convex, where J is the normalized duality map (see definition 2.1). The variational inequality problem, which has its origin in the 1964 result of Stampacchia [21], has engaged the interest of researchers in the recent past (see, e.g., [25, 26] and many others). This is concerned with the following: For a monotone operator $A : C \rightarrow E$, find a point $x^* \in C$ such that

$$(1.1) \quad \langle y - x^*, Ax^* \rangle \geq 0 \text{ for all } y \in C.$$

The set of solutions of (1.1) is denoted by $VI(C, A)$. This problem, which plays a crucial role in nonlinear analysis, is also related to fixed point problems, zeros of nonlinear operators, complementarity problems, and convex minimization problems (see, for example, [29, 30]).

A related problem is the equilibrium problem, which has been studied by several researchers and is mostly applied in solving optimization problems (see [3]). For a map $f : C \rightarrow E$, the equilibrium problem is concerned with finding a point $x^* \in C$ such that

$$(1.2) \quad f(x^*, y) \geq 0 \text{ for all } y \in C.$$

The set of solutions of (1.2) is denoted by $EP(f)$. The variational inequality and equilibrium problems are special cases of the so-called generalized mixed equilibrium problem (see [16]). Another related problem is the fixed point problem. For a map $T : D(T) \subset E \rightarrow E$, the fixed points of T are the points $x^* \in D(T)$ such that $Tx^* = x^*$. Recently, owing to the need to develop methods for solving fixed points of problems for functions

Received: 22.11.2021. In revised form: 07.07.2022. Accepted: 14.07.2022

2010 *Mathematics Subject Classification.* 47H09, 47H05, 47J25, 47J05.

Key words and phrases. *equilibrium problem, J_* -nonexpansive, fixed points, variational inequality, strong convergence.*

Corresponding author: Maria A. Onyido; mao0021@auburn.edu

from a space to its dual, a new concept of *fixed points for maps from a real normed space E to its dual space E^** , called J -fixed point has been introduced and studied (see [6, 13, 24]). With this evolving fixed point theory, we study the J -fixed points of certain maps and the following equilibrium problem. Let $f : JC \times JC \rightarrow \mathbb{R}$ be a bifunction. The equilibrium problem for f is finding

$$(1.3) \quad x^* \in C \text{ such that } f(Jx^*, Jy) \geq 0, \forall y \in C.$$

We denote the solution set of (1.3) by $EP(f)$. Several problems in physics, optimization and economics reduce to finding a solution of (1.3) (see, e.g., [8, 25] and the references in them). Most of the equilibrium problems studied in the past two decades centered on their existence and applications (see, e.g., [3, 8]). However, recently, several researchers have started working on finding approximate solutions of equilibrium problems and their generalizations (see, e.g., [12, 26]). Not long ago, some researchers investigated the problem of establishing a common element in the solution set of an equilibrium problem, fixed point of a family of nonexpansive maps and solution set of a variational inequality problem for different classes of maps (see [27] and references therein).

In this paper, inspired by the above results especially the works in [4, 23, 27], we present an algorithm for finding a common element of the fixed point of an infinite family of generalized J_* -nonexpansive maps, the solution set of the variational inequality problem of a finite family of continuous monotone maps and the solution set of the equilibrium point of a finite family of bifunctions satisfying some given conditions. Our results complement, generalize and extend results in [14, 19, 20, 27] (see the section on conclusion) and other recent results in this direction. It is worth noting that very recently, the authors in [4] introduced a new class of maps which they called *relatively weak J -nonexpansive* and developed an algorithm for approximating a common element of the J -fixed point of a countable family of such maps and zeros of some other class of maps in certain Banach spaces. Previously, maps with similar requirements as these *relatively weak J -nonexpansive* maps have also been studied in [7] where they were called *quasi- ϕ - J -nonexpansive*. We observe that these two sets of maps (*relatively weak J -nonexpansive* and *quasi- ϕ - J -nonexpansive*) coincide in definition with the J_* -nonexpansive maps in our results.

2. PRELIMINARIES

In this section, we present definitions and lemmas used in proving our main results.

Definition 2.1. (Normalized duality map) The map $J : E \rightarrow 2^{E^*}$ defined by

$$Jx := \{x^* \in E^* : \langle x, x^* \rangle = \|x\| \cdot \|x^*\|, \|x\| = \|x^*\|\}$$

is called the *normalized duality map* on E .

It is well known that if E is smooth, strictly convex and reflexive then J^{-1} exists (see e.g., [11]); $J^{-1} : E^* \rightarrow E$ is the normalized duality mapping on E^* , and $J^{-1} = J_*$, $JJ_* = I_{E^*}$ and $J_*J = I_E$, where I_E and I_{E^*} are the identity maps on E and E^* , respectively. A well known property of J is, see e.g., [9, 11], if E is uniformly smooth, then J is uniformly continuous on bounded subsets of E .

Definition 2.2. (Lyapunov Functional) [1, 12] Let E be a smooth real Banach space with dual E^* . The *Lyapunov functional* $\phi : E \times E \rightarrow \mathbb{R}$, is defined by

$$(2.4) \quad \phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2, \text{ for } x, y \in E,$$

where J is the normalized duality map. If $E = H$, a real Hilbert space, then equation (2.4) reduces to $\phi(x, y) = \|x - y\|^2$ for $x, y \in H$. Additionally,

$$(2.5) \quad (\|x\| - \|y\|)^2 \leq \phi(x, y) \leq (\|x\| + \|y\|)^2 \text{ for } x, y \in E.$$

Definition 2.3. (Generalized nonexpansive) [17, 18] Let C be a nonempty closed and convex subset of a real Banach space E and T be a map from C to E . The map T is called *generalized nonexpansive* if $F(T) := \{x \in C : Tx = x\} \neq \emptyset$ and $\phi(Tx, p) \leq \phi(x, p)$ for all $x \in C, p \in F(T)$.

Definition 2.4. (Retraction) [17, 18] A map R from E onto C is said to be a retraction if $R^2 = R$. The map R is said to be *sunny* if $R(Rx + t(x - Rx)) = Rx$ for all $x \in E$ and $t \leq 0$.

A nonempty closed subset C of a smooth Banach space E is said to be a *sunny generalized nonexpansive retract* of E if there exists a sunny generalized nonexpansive retraction R from E onto C .

NST-condition. Let C be a closed subset of a Banach space E . Let $\{T_n\}$ and Γ be two families of generalized nonexpansive maps of C into E such that $\bigcap_{n=1}^{\infty} F(T_n) = F(\Gamma) \neq \emptyset$, where $F(T_n)$ is the set of fixed points of $\{T_n\}$ and $F(\Gamma)$ is the set of common fixed points of Γ .

Definition 2.5. [17] The sequence $\{T_n\}$ satisfies the NST-condition (see e.g., [15]) with Γ if for each bounded sequence $\{x_n\} \subset C$,

$$\lim_{n \rightarrow \infty} \|x_n - T_n x_n\| = 0 \Rightarrow \lim_{n \rightarrow \infty} \|x_n - T x_n\| = 0, \text{ for all } T \in \Gamma.$$

Remark 2.1. If $\Gamma = \{T\}$ a singleton, $\{T_n\}$ satisfies the NST-condition with $\{T\}$. If $T_n = T$ for all $n \geq 1$, then, $\{T_n\}$ satisfies the NST-condition with $\{T\}$.

Let C be a nonempty closed and convex subset of a uniformly smooth and uniformly convex real Banach space E with dual space E^* . Let J be the normalized duality map on E and J_* be the normalized duality map on E^* . Observe that under this setting, J^{-1} exists and $J^{-1} = J_*$. With these notations, we have the following definitions.

Definition 2.6. (Closed map) [23] A map $T : C \rightarrow E^*$ is called J_* -closed if $(J_* \circ T) : C \rightarrow E$ is a closed map, i.e., if $\{x_n\}$ is a sequence in C such that $x_n \rightarrow x$ and $(J_* \circ T)x_n \rightarrow y$, then $(J_* \circ T)x = y$.

Definition 2.7. (J -fixed Point) [6] A point $x^* \in C$ is called a J -fixed point of T if $Tx^* = Jx^*$. The set of J -fixed points of T will be denoted by $F_J(T)$.

Definition 2.8. (Generalized J_* -nonexpansive) [23] A map $T : C \rightarrow E^*$ will be called *generalized J_* -nonexpansive* if $F_J(T) \neq \emptyset$, and $\phi(p, (J_* \circ T)x) \leq \phi(p, x)$ for all $x \in C$ and for all $p \in F_J(T)$.

Remark 2.2. Examples of generalized J_* -nonexpansive maps in Hilbert and more general Banach spaces were given in [4, 23].

Let C be a nonempty closed subset of a smooth, strictly convex and reflexive Banach space E such that JC is closed and convex. For solving the equilibrium problem, let us assume that a bifunction $f : JC \times JC \rightarrow \mathbb{R}$ satisfies the following conditions:

- (A1) $f(x^*, x^*) = 0$ for all $x^* \in JC$;
- (A2) f is monotone, i.e. $f(x^*, y^*) + f(y^*, x^*) \leq 0$ for all $x^*, y^* \in JC$;
- (A3) for all $x^*, y^*, z^* \in JC$, $\limsup_{t \downarrow 0} f(tz^* + (1-t)x^*, y^*) \leq f(x^*, y^*)$;
- (A4) for all $x^* \in JC$, $f(x^*, \cdot)$ is convex and lower semicontinuous.

With the above definitions, we now provide the lemmas we shall use.

Lemma 2.1. [28] Let E be a uniformly convex Banach space, $r > 0$ be a positive number, and $B_r(0)$ be a closed ball of E . For any given points $\{x_1, x_2, \dots, x_N\} \subset B_r(0)$ and any given positive numbers $\{\lambda_1, \lambda_2, \dots, \lambda_N\}$ with $\sum_{n=1}^N \lambda_n = 1$, there exists a continuous strictly increasing and convex function $g : [0, 2r) \rightarrow [0, \infty)$ with $g(0) = 0$ such that, for any $i, j \in \{1, 2, \dots, N\}$, $i < j$,

$$(2.6) \quad \left\| \sum_{n=1}^N \lambda_n x_n \right\|^2 \leq \sum_{n=1}^N \lambda_n \|x_n\|^2 - \lambda_i \lambda_j g(\|x_i - x_j\|).$$

Lemma 2.2. [12] Let X be a real smooth and uniformly convex Banach space, and let $\{x_n\}$ and $\{y_n\}$ be two sequences of X . If either $\{x_n\}$ or $\{y_n\}$ is bounded and $\phi(x_n, y_n) \rightarrow 0$ as $n \rightarrow \infty$, then $\|x_n - y_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 2.3. [1] Let C be a nonempty closed and convex subset of a smooth, strictly convex and reflexive Banach space E . Then, the following are equivalent.

- (i) C is a sunny generalized nonexpansive retract of E ,
- (ii) C is a generalized nonexpansive retract of E ,
- (iii) JC is closed and convex.

Lemma 2.4. [1] Let C be a nonempty closed and convex subset of a smooth and strictly convex Banach space E such that there exists a sunny generalized nonexpansive retraction R from E onto C . Then, the following hold.

- (i) $z = Rx$ iff $\langle x - z, Jy - Jz \rangle \leq 0$ for all $y \in C$,
- (ii) $\phi(x, Rx) + \phi(Rx, z) \leq \phi(x, z)$.

Lemma 2.5. [10] Let C be a nonempty closed sunny generalized nonexpansive retract of a smooth and strictly convex Banach space E . Then the sunny generalized nonexpansive retraction from E to C is uniquely determined.

Lemma 2.6. [3] Let C be a nonempty closed subset of a smooth, strictly convex and reflexive Banach space E such that JC is closed and convex, let f be a bifunction from $JC \times JC$ to \mathbb{R} satisfying (A1) – (A4). For $r > 0$ and let $x \in E$. Then there exists $z \in C$ such that $f(Jz, Jy) + \frac{1}{r} \langle z - x, Jy - Jz \rangle \geq 0, \forall y \in C$.

Lemma 2.7. [22] Let C be a nonempty closed subset of a smooth, strictly convex and reflexive Banach space E such that JC is closed and convex, let f be a bifunction from $JC \times JC$ to \mathbb{R} satisfying (A1) – (A4). For $r > 0$ and let $x \in E$, define a mapping $T_r(x) : E \rightarrow C$ as follows:

$$T_r(x) = \{z \in C : f(Jz, Jy) + \frac{1}{r} \langle y - z, Jz - Jx \rangle \geq 0, \forall y \in C\}.$$

Then the following hold:

- (i) T_r is single valued;
- (ii) for all $x, y \in E, \langle T_r x - T_r y, JT_r x - JT_r y \rangle \leq \langle x - y, JT_r x - JT_r y \rangle$;
- (iii) $F(T_r) = EP(f)$;
- (iv) $\phi(p, T_r(x)) + \phi(T_r(x), x) \leq \phi(p, x)$ for all $p \in F(T_r)$.
- (v) $JEP(f)$ is closed and convex.

Lemma 2.8. [23] Let C be a nonempty closed subset of a smooth, strictly convex and reflexive Banach space E . Let $A : C \rightarrow E^*$ be a continuous monotone mapping. For $r > 0$ and let $x \in E$, define a mapping $F_r(x) : E \rightarrow C$ as follows:

$$F_r(x) = \{z \in C : \langle y - z, Az \rangle + \frac{1}{r} \langle y - z, Jz - Jx \rangle \geq 0, \forall y \in C\}.$$

Then the following hold:

- (i) F_r is single valued;
- (ii) for all $x, y \in E$, $\langle F_r x - T_r y, JF_r x - JF_r y \rangle \leq \langle x - y, JF_r x - JF_r y \rangle$;
- (iii) $F(F_r) = VI(C, A)$;
- (iv) $\phi(p, F_r(x)) + \phi(F_r(x), x) \leq \phi(p, x)$ for all $p \in F(F_r)$.
- (v) $JVI(C, A)$ is closed and convex.

Lemma 2.9. [23] Let E be a uniformly convex and uniformly smooth real Banach space with dual space E^* and let C be a closed subset of E such that JC is closed and convex. Let T be a generalized J_* -nonexpansive map from C to E^* such that $F_J(T) \neq \emptyset$, then $F_J(T)$ and $JF_J(T)$ are closed. Additionally, if $JF_J(T)$ is convex, then $F_J(T)$ is a sunny generalized nonexpansive retract of E .

3. MAIN RESULTS

Let E be a uniformly smooth and uniformly convex real Banach space with dual space E^* and let C be a nonempty closed and convex subset of E such that JC is closed and convex. Let $f_l, l = 1, 2, 3, \dots, L$ be a family of bifunctions from $JC \times JC$ to \mathbb{R} satisfying (A1) – (A4), $T_n : C \rightarrow E^*, n = 1, 2, 3, \dots$ be an infinite family of generalized J_* -nonexpansive maps, and $A_k : C \rightarrow E^*, k = 1, 2, 3, \dots, N$ be a finite family of continuous monotone mappings. Let the sequence $\{x_n\}$ be generated by the following iteration process:

$$(3.7) \quad \begin{cases} x_1 = x \in C; C_1 = C, \\ z_n := \{z \in C : f_n(Jz, Jy) + \frac{1}{r_n} \langle y - z, Jz - Jx_n \rangle \geq 0, \forall y \in C\}, \\ u_n := \{z \in C : \langle y - z, A_n z \rangle + \frac{1}{r_n} \langle y - z, Jz - Jx_n \rangle \geq 0, \forall y \in C\}, \\ y_n = J^{-1}(\alpha_1 Jx_n + \alpha_2 Jz_n + \alpha_3 T_n u_n), \\ C_{n+1} = \{z \in C_n : \phi(z, y_n) \leq \phi(z, x_n)\}, \\ x_{n+1} = R_{C_{n+1}} x, \end{cases}$$

for all $n \in \mathbb{N}$, with $\alpha_1, \alpha_2, \alpha_3 \in (0, 1)$ satisfying $\alpha_1 + \alpha_2 + \alpha_3 = 1$, $\{r_n\} \subset [a, \infty)$ for some $a > 0$, $A_n = A_{n(\text{mod } N)}$ and $f_n(\cdot, \cdot) = f_{n(\text{mod } L)}(\cdot, \cdot)$.

Lemma 3.10. The sequence $\{x_n\}$ generated by (3.7) is well defined.

Proof. Observe that JC_1 is closed and convex. Moreover, it is easy to see that $\phi(z, y_n) \leq \phi(z, x_n)$ is equivalent to

$$0 \leq \|x_n\|^2 - \|y_n\|^2 - 2\langle z, Jx_n - Jy_n \rangle,$$

which is affine in z . Hence, by induction JC_n is closed and convex for each $n \geq 1$. Therefore, from Lemma 2.3, we have that C_n is a sunny generalized retract of E for each $n \geq 1$. This shows that $\{x_n\}$ is well defined. \square

Theorem 3.1. Let E be a uniformly smooth and uniformly convex real Banach space with dual space E^* and let C be a nonempty closed and convex subset of E such that JC is closed and convex. Let $f_l, l = 1, 2, 3, \dots, L$ be a family of bifunctions from $JC \times JC$ to \mathbb{R} satisfying (A1) – (A4), $T_n : C \rightarrow E^*, n = 1, 2, 3, \dots$ be an infinite family of generalized J_* -nonexpansive maps, $A_k : C \rightarrow E^*, k = 1, 2, 3, \dots, N$ be a finite family of continuous monotone mappings and Γ be a family of J_* -closed and generalized J_* -nonexpansive maps from C to E^* such that $\bigcap_{n=1}^{\infty} F_J(T_n) = F_J(\Gamma) \neq \emptyset$ and $B := F_J(\Gamma) \cap \left[\bigcap_{l=1}^L EP(f_l) \right] \cap \left[\bigcap_{k=1}^N VI(C, A_k) \right] \neq \emptyset$. Assume that $JF_J(\Gamma)$ is convex and $\{T_n\}$ satisfies the NST-condition with Γ . Then, $\{x_n\}$ generated by (3.7) converges strongly to $R_B x$, where R_B is the sunny generalized nonexpansive retraction of E onto B .

Proof. The proof is given in 6 steps.

Step 1: We show that the expected limit R_Bx exists as a point in C_n for all $n \geq 1$.

First, we show that $B \subset C_n$ for all $n \geq 1$ and B is a sunny generalized retract of E .

Since $C_1 = C$, we have $B \subset C_1$. Suppose $B \subset C_n$ for some $n \in \mathbb{N}$. Let $u \in B$. We observe from algorithm (3.7) that $u_n = F_{r_n}x_n$ and $z_n = T_{r_n}x_n$ for all $n \in \mathbb{N}$, using this and the fact that $\{T_n\}$ is an infinite family of generalized J_* -nonexpansive maps, the definition of y_n , Lemmas 2.7, 2.8, and 2.1, we compute as follows:

$$\begin{aligned}
 \phi(u, y_n) &= \phi(u, J^{-1}(\alpha_1 Jx_n + \alpha_2 Jz_n + \alpha_3 T_n u_n)) \\
 &\leq \alpha_1 [\|u\|^2 - 2\langle u, Jx_n \rangle + \|x_n\|^2] + \alpha_2 [\|u\|^2 - 2\langle u, Jz_n \rangle + \|z_n\|^2] \\
 &\quad + \alpha_3 [\|u\|^2 - 2\langle u, J(J_* \circ T_n)u_n \rangle + \|T_n u_n\|^2] \\
 &\quad - \alpha_1 \alpha_3 g(\|Jx_n - J(J_* \circ T_n)u_n\|) \\
 &= \alpha_1 \phi(u, x_n) + \alpha_2 \phi(u, z_n) + \alpha_3 \phi(u, (J_* \circ T_n)u_n) - \alpha_1 \alpha_3 g(\|Jx_n - T_n u_n\|) \\
 (3.8) \quad &\leq \alpha_1 \phi(u, x_n) + \alpha_2 \phi(u, z_n) + \alpha_3 \phi(u, u_n) - \alpha_1 \alpha_3 g(\|Jx_n - T_n u_n\|) \\
 &= \alpha_1 \phi(u, x_n) + \alpha_2 \phi(u, T_{r_n}x_n) + \alpha_3 \phi(u, u_n) - \alpha_1 \alpha_3 g(\|Jx_n - T_n u_n\|) \\
 &\leq \alpha_1 \phi(u, x_n) + \alpha_2 \phi(u, x_n) + \alpha_3 \phi(u, u_n) - \alpha_1 \alpha_3 g(\|Jx_n - T_n u_n\|),
 \end{aligned}$$

which yields

$$(3.9) \quad \phi(u, y_n) \leq \phi(u, x_n) - \alpha_1 \alpha_3 g(\|Jx_n - T_n u_n\|).$$

Hence, $\phi(u, y_n) \leq \phi(u, x_n)$ and we have that $u \in C_{n+1}$, which implies that $B \subset C_n$ for all $n \geq 1$. Moreover, From Lemma 2.7 and 2.8 both $JVI(C, A_k)$ and $JEP(f_l)$ are closed and convex for each l and for each k . Also, using our assumption and lemma 2.9, we have that $J(F_J(\Gamma))$ is closed and convex. Since E is uniformly convex, J is one-to-one. Thus, we have that,

$$J\left(F_J(\Gamma) \cap \left[\bigcap_{l=1}^L EP(f_l) \right] \cap \left[\bigcap_{k=1}^N VI(C, A_k) \right] \right) = JF_J(\Gamma) \cap J\left[\bigcap_{l=1}^L EP(f_l) \right] \cap J\left[\bigcap_{k=1}^N VI(C, A_k) \right]$$

so $J(B)$ is closed and convex. Using Lemma 2.3, we obtain that B is a sunny generalized retract of E . Therefore, from Lemma 2.5, we have that R_Bx exists as a point in C_n for all $n \geq 1$. This completes step 1.

Step 2: We show that the sequence $\{x_n\}$ defined by (3.7) converges to some $x^* \in C$. Using the fact that $x_n = R_{C_n}x$ and Lemma 2.4(ii), we obtain

$$\phi(x, x_n) = \phi(x, R_{C_n}x) \leq \phi(x, u) - \phi(R_{C_n}x, u) \leq \phi(x, u),$$

for all $u \in F_J(\Gamma) \cap EP(f_l) \cap VI(C, A_k) \subset C_n$; ($l = 1, 2, \dots, L$; $k = 1, 2, \dots, K$). This implies that $\{\phi(x, x_n)\}$ is bounded. Hence, from equation (2.5), $\{x_n\}$ is bounded. Also, since $x_{n+1} = R_{C_{n+1}}x \in C_{n+1} \subset C_n$, and $x_n = R_{C_n}x \in C_n$, applying Lemma 2.4(ii) gives

$$\phi(x, x_n) \leq \phi(x, x_{n+1}) \quad \forall n \in \mathbb{N}.$$

So, $\lim_{n \rightarrow \infty} \phi(x, x_n)$ exists. Again, using Lemma 2.4(ii) and $x_n = R_{C_n}x$, we obtain that for all $m, n \in \mathbb{N}$ with $m > n$,

$$\begin{aligned}
 \phi(x_n, x_m) &= \phi(R_{C_n}x, x_m) \leq \phi(x, x_m) - \phi(x, R_{C_n}x) \\
 (3.10) \quad &= \phi(x, x_m) - \phi(x, x_n) \rightarrow 0 \text{ as } n \rightarrow \infty.
 \end{aligned}$$

From Lemma 2.2, we conclude that $\|x_n - x_m\| \rightarrow 0$, as $m, n \rightarrow \infty$. Hence, $\{x_n\}$ is a Cauchy sequence in C , and so, there exists $x^* \in C$ such that $x_n \rightarrow x^*$ completing step 2.

Step 3: We prove $x^* \in \bigcap_{k=1}^N VI(C, A_k)$.

From the definitions of C_{n+1} and x_{n+1} , we obtain that $\phi(x_{n+1}, y_n) \leq \phi(x_{n+1}, x_n) \rightarrow 0$ as $n \rightarrow \infty$. Hence, by Lemma 2.2, we have that

$$(3.11) \quad \lim_{n \rightarrow \infty} \|x_n - y_n\| = 0.$$

Since from step 2 $x_n \rightarrow x^*$ as $n \rightarrow \infty$, equation (3.11) implies that $y_n \rightarrow x^*$ as $n \rightarrow \infty$. Using the fact that $u_n = F_{r_n} x_n$ for all $n \in \mathbb{N}$ and Lemma 2.2, we get for $u \in B$,

$$(3.12) \quad \begin{aligned} \phi(u_n, x_n) &= \phi(F_{r_n} x_n, x_n) \\ &\leq \phi(u, x_n) - \phi(u, F_{r_n} x_n) \\ &= \phi(u, x_n) - \phi(u, u_n). \end{aligned}$$

From equations (3.8) and (3.9) we have

$$(3.13) \quad \phi(u, y_n) \leq \alpha_1 \phi(u, x_n) + \alpha_2 \phi(u, x_n) + \alpha_3 \phi(u, u_n) \leq \phi(u, x_n).$$

Since $x_n, y_n \rightarrow x^*$ as $n \rightarrow \infty$, equation (3.13) implies that $\phi(u, u_n) \rightarrow \phi(u, x^*)$ as $n \rightarrow \infty$. Therefore, from (3.12), we have $\phi(u, x_n) - \phi(u, u_n) \rightarrow 0$ as $n \rightarrow \infty$ which implies that $\lim_{n \rightarrow \infty} \phi(u_n, x_n) = 0$. Hence, from Lemma 2.2, we have

$$(3.14) \quad \lim_{n \rightarrow \infty} \|u_n - x_n\| = 0.$$

Observe that since J is uniformly continuous on bounded subsets of E , it follows from (3.14) that $\|Ju_n - Jx_n\| \rightarrow 0$.

Again, since $r_n \in [a, \infty)$, we have that

$$(3.15) \quad \lim_{n \rightarrow \infty} \frac{\|Ju_n - Jx_n\|}{r_n} = 0.$$

From $u_n = F_{r_n} x_n$, we have

$$(3.16) \quad \langle y - u_n, A_n u_n \rangle + \frac{1}{r_n} \langle y - u_n, Ju_n - Jx_n \rangle \geq 0, \quad \forall y \in C.$$

Let $\{n_l\}_{l=1}^{\infty} \subset \mathbb{N}$ be such that $A_{n_l} = A_1 \forall l \geq 1$. Then, from (3.16), we obtain

$$(3.17) \quad \langle y - u_{n_l}, A_1 u_{n_l} \rangle + \frac{1}{r_{n_l}} \langle y - u_{n_l}, Ju_{n_l} - Jx_{n_l} \rangle \geq 0, \quad \forall y \in C.$$

If we set $v_t = ty + (1-t)x^*$ for all $t \in (0, 1]$ and $y \in C$, then we get that $v_t \in C$. Hence, it follows from (3.17) that

$$(3.18) \quad \langle v_t - u_{n_l}, A_1 u_{n_l} \rangle + \langle y - u_{n_l}, \frac{Ju_{n_l} - Jx_{n_l}}{r_{n_l}} \rangle \geq 0.$$

This implies that

$$\begin{aligned} \langle v_t - u_{n_l}, A_1 v_t \rangle &\geq \langle v_t - u_{n_l}, A_1 v_t \rangle - \langle v_t - u_{n_l}, A_1 u_{n_l} \rangle - \langle y - u_{n_l}, \frac{Ju_{n_l} - Jx_{n_l}}{r_{n_l}} \rangle \\ &= \langle v_t - u_{n_l}, A_1 v_t - A_1 u_{n_l} \rangle - \langle y - u_{n_l}, \frac{Ju_{n_l} - Jx_{n_l}}{r_{n_l}} \rangle. \end{aligned}$$

Since A_1 is monotone, $\langle v_t - u_{n_l}, A_1 v_t - A_1 u_{n_l} \rangle \geq 0$. Thus, using (3.15), we have that

$$0 \leq \lim_{l \rightarrow \infty} \langle v_t - u_{n_l}, A_1 v_t \rangle = \langle v_t - x^*, A_1 v_t \rangle,$$

therefore,

$$\langle y - x^*, A_1 v_t \rangle \geq 0, \quad \forall y \in C.$$

Letting $t \rightarrow 0$ and using continuity of A_1 , we have that

$$\langle y - x^*, A_1 x^* \rangle \geq 0, \quad \forall y \in C.$$

This implies that $x^* \in VI(C, A_1)$. Similarly, if $\{n_i\}_{i=1}^{\infty} \subset \mathbb{N}$ is such that $A_{n_i} = A_2$ for all $i \geq 1$, then we have again that $x^* \in VI(C, A_2)$. If we continue in similar manner, we obtain that $x^* \in \bigcap_{k=1}^N VI(C, A_k)$.

Step 4: We prove that $x^* \in F_J(\Gamma)$.

First, we show that $\lim_{n \rightarrow \infty} \|Jx_n - Tu_n\| = 0 \forall T \in \Gamma$.

From inequality (3.9) and the fact that g is nonnegative, we obtain

$$0 \leq \alpha_1 \alpha_3 g(\|Jx_n - Tu_n\|) \leq \phi(u, x_n) - \phi(u, y_n) \leq 2\|u\| \cdot \|Jx_n - Jy_n\| + \|x_n - y_n\|M,$$

for some $M > 0$. Thus, using (3.11) and properties of g , we obtain that

$\lim_{n \rightarrow \infty} \|Jx_n - Tu_n\| = 0$. Using the above and triangle inequality gives $\|Ju_n - Tu_n\| \rightarrow 0$ as $n \rightarrow \infty$. Since $\{T_n\}_{n=1}^\infty$ satisfies the NST condition with Γ , we have that

$$(3.19) \quad \lim_{n \rightarrow \infty} \|Ju_n - Tu_n\| = 0 \forall T \in \Gamma.$$

Now, from equation (3.14), we have $u_n \rightarrow x^* \in C$. Assume that $(J_* \circ T)u_n \rightarrow y^*$. Since T is J_* -closed, we have $y^* = (J_* \circ T)x^*$. Furthermore, by the uniform continuity of J on bounded subsets of E , we have: $Ju_n \rightarrow Jx^*$ and $J(J_* \circ T)u_n \rightarrow Jy^*$ as $n \rightarrow \infty$. Hence, we have

$$\lim_{n \rightarrow \infty} \|Ju_n - J(J_* \circ T)u_n\| = \lim_{n \rightarrow \infty} \|Ju_n - Tu_n\| = 0, \forall T \in \Gamma,$$

which implies $\|Jx^* - Jy^*\| = \|Jx^* - J(J_* \circ T)x^*\| = \|Jx^* - Tx^*\| = 0$. So, $x^* \in F_J(\Gamma)$.

Step 5: We prove that $x^* \in \cap_{l=1}^L EP(f_l)$.

This follows by similar argument as in step 3 but for the sake of completeness we provide the details. Using the fact that $z_n = T_{r_n}x_n$ and Lemma 2.7, we obtain that for $u \in F_J(\Gamma) \cap EP(f_l) \cap VI(C, A_k)$ for all i, k ,

$$(3.20) \quad \begin{aligned} \phi(z_n, x_n) &= \phi(T_{r_n}x_n, x_n) \\ &\leq \phi(u, x_n) - \phi(u, T_{r_n}x_n) \\ &= \phi(u, x_n) - \phi(u, z_n). \end{aligned}$$

From equations (3.8) and (3.9), we have

$$(3.21) \quad \phi(u, y_n) \leq \alpha_1 \phi(u, x_n) + \alpha_2 \phi(u, z_n) + \alpha_3 \phi(u, x_n) \leq \phi(u, x_n).$$

Since $x_n, y_n, u_n \rightarrow x^*$ as $n \rightarrow \infty$, from equation (3.21) we have $\phi(u, z_n) \rightarrow \phi(u, x^*)$ as $n \rightarrow \infty$. Therefore, from (3.20), we have $\phi(u, x_n) - \phi(u, u_n) \rightarrow 0$ as $n \rightarrow \infty$. Hence $\lim_{n \rightarrow \infty} \phi(z_n, x_n) = 0$. From Lemma 2.2, we have

$$(3.22) \quad \lim_{n \rightarrow \infty} \|z_n - x_n\| = 0,$$

which implies that $z_n \rightarrow x^*$ as $n \rightarrow \infty$. Again, since J is uniformly continuous on bounded subsets of E , (3.22) implies $\|Jz_n - Jx_n\| \rightarrow 0$. Since $r_n \in [a, \infty)$, we have that

$$(3.23) \quad \lim_{n \rightarrow \infty} \frac{\|Jz_n - Jx_n\|}{r_n} = 0.$$

Since $z_n = T_{r_n}x_n$, we have that

$$\frac{1}{r_n} \langle y - z_n, Jz_n - Jx_n \rangle \geq -f_n(Jz_n, Jy), \forall y \in C.$$

Let $\{n_l\}_{l=1}^\infty \subset \mathbb{N}$ be such that $f_{n_l} = f_1 \forall l \geq 1$. Then, using (A2), we have

$$(3.24) \quad \langle y - z_n, \frac{Jz_n - Jx_n}{r_n} \rangle \geq -f_1(Jz_n, Jy) \geq f_1(Jy, Jz_n), \forall y \in C.$$

Since $f_1(x, \cdot)$ is convex and lower-semicontinuous and $z_n \rightarrow x^*$, it follows from equation (3.23) and inequality (3.24) that

$$f_1(Jy, Jx^*) \leq 0, \forall y \in C.$$

For $t \in (0, 1]$ and $y \in C$, let $y_t^* = tJy + (1 - t)Jx^*$. Since JC is convex, we have that $y_t^* \in JC$ and hence $f_1(y_t^*, Jx^*) \leq 0$. Applying (A1) gives,

$$0 = f_1(y_t^*, y_t^*) \leq tf_1(y_t^*, Jy) + (1 - t)f_1(y_t^*, Jx^*) \leq tf_1(y_t^*, Jy), \forall y \in C.$$

This implies that

$$f_1(y_t^*, Jy) \geq 0, \forall y \in C.$$

Letting $t \downarrow 0$ and using (A3), we get

$$f_1(Jx^*, Jy) \geq 0, \forall y \in C.$$

Therefore, we have that $Jx^* \in JEP(f_1)$. This implies that $x^* \in EP(f_1)$. Applying similar argument, we can show that $x^* \in EP(f_l)$ for $l = 2, 3, \dots, L$. Hence, $x^* \in \bigcap_{l=1}^L EP(f_l)$.

Step 6: Finally, we show that $x^* = R_Bx$.

From Lemma 2.4(ii), we obtain that

$$(3.25) \quad \phi(x, R_Bx) \leq \phi(x, x^*) - \phi(R_Bx, x^*) \leq \phi(x, x^*).$$

Again, using Lemma 2.4(ii), definition of x_{n+1} , and $x^* \in B \subset C_n$, we compute as follows:

$$\begin{aligned} \phi(x, x_{n+1}) &\leq \phi(x, x_{n+1}) + \phi(x_{n+1}, R_Bx) \\ &= \phi(x, R_{C_{n+1}}x) + \phi(R_{C_{n+1}}x, R_Bx) \leq \phi(x, R_Bx). \end{aligned}$$

Since $x_n \rightarrow x^*$, taking limits on both sides of the last inequality, we obtain

$$(3.26) \quad \phi(x, x^*) \leq \phi(x, R_Bx).$$

Using inequalities (3.25) and (3.26), we obtain that $\phi(x, x^*) = \phi(x, R_Bx)$. By the uniqueness of R_B (Lemma 2.5), we obtain that $x^* = R_Bx$. This completes proof of the theorem. \square

4. APPLICATIONS

Corollary 4.1. *Let E be a uniformly smooth and uniformly convex real Banach space with dual space E^* and let C be a nonempty closed and convex subset of E such that JC is closed and convex. Let f be a bifunction from $JC \times JC$ to \mathbb{R} satisfying (A1) – (A4), $A : C \rightarrow E^*$, be a continuous monotone mapping, $T : C \rightarrow E^*$, be a generalized J_* -nonexpansive and J_* -closed map such that $B := F_J(T) \cap EP(f) \cap VI(C, A) \neq \emptyset$. Assume that $JF_J(T)$ is convex. Then, $\{x_n\}$ generated by (3.7) converges strongly to R_Bx , where R_B is the sunny generalized nonexpansive retraction of E onto B .*

Proof. Set $T_n := T$ for all $n \in \mathbb{N}$, $A := A_i$ for any $i = 1, 2, \dots, N$, and $f := f_l$ for any $l = 1, 2, \dots, L$. Then, from remark 2.1, $\{T_n\}$ satisfies the NST-condition with $\{T\}$. The conclusion follows from Theorem 3.1. \square

Corollary 4.2. *Let E be a uniformly smooth and uniformly convex real Banach space with dual space E^* and let C be a nonempty closed and convex subset of E such that JC is closed and convex. Let $f_l, l = 1, 2, 3, \dots, L$ be a family of bifunctions from $JC \times JC$ to \mathbb{R} satisfying (A1) – (A4), $T_n : C \rightarrow E^*, n = 1, 2, 3, \dots$ be an infinite family of generalized J_* -nonexpansive maps and Γ be a family of J_* -closed and generalized J_* -nonexpansive maps from C to E^* such that $\bigcap_{n=1}^\infty F_J(T_n) = F_J(\Gamma) \neq \emptyset$ and $B := F_J(\Gamma) \cap \left[\bigcap_{l=1}^L EP(f_l) \right] \neq \emptyset$. Assume that $JF_J(\Gamma)$ is convex and $\{T_n\}$ satisfies the NST-condition with Γ . Then, $\{x_n\}$ generated by (3.7) converges strongly to R_Bx , where R_B is the sunny generalized nonexpansive retraction of E onto B .*

Proof. Setting $A_k = 0$ for any $k = 1, 2, 3, \dots, N$, then result follows from Theorem 3.1. \square

Remark 4.3. We note here that the theorem and corollaries presented above are applicable in classical Banach spaces, such as L_p , l_p , or $W_p^m(\Omega)$, $1 < p < \infty$, where $W_p^m(\Omega)$ denotes the usual Sobolev space.

Remark 4.4. ([2]; p. 36) The analytical representations of duality maps are known in a number of Banach spaces, for example, in the spaces L_p , l_p , and $W_m^p(\Omega)$, $p \in (1, \infty)$, $p^{-1} + q^{-1} = 1$.

Corollary 4.3. Let $E = H$, a real Hilbert space and let C be a nonempty closed and convex subset of H . Let $f_l, l = 1, 2, 3, \dots, L$ be a family of bifunctions from $C \times C$ to \mathbb{R} satisfying (A1) – (A4), $T_n : C \rightarrow H, n = 1, 2, 3, \dots$ be an infinite family of nonexpansive maps, $A_k : C \rightarrow H, k = 1, 2, 3, \dots, N$ be a finite family of continuous monotone mappings and Γ be a family of closed and generalized nonexpansive maps from C to H such that $\bigcap_{n=1}^\infty F(T_n) = F(\Gamma) \neq \emptyset$ and $B := F(\Gamma) \cap \left[\bigcap_{l=1}^L EP(f_l) \right] \cap \left[\bigcap_{k=1}^N VI(C, A_k) \right] \neq \emptyset$. Assume that $\{T_n\}$ satisfies the NST-condition with Γ . Let $\{x_n\}$ be generated by:

$$(4.27) \quad \begin{cases} x_1 = x \in C; C_1 = C, \\ z_n := \{z \in C : f_n(z, y) + \frac{1}{r_n} \langle y - z, z - x_n \rangle \geq 0, \forall y \in C\}, \\ u_n := \{z \in C : \langle y - z, A_n z \rangle + \frac{1}{r_n} \langle y - z, z - x_n \rangle \geq 0, \forall y \in C\}, \\ y_n = \alpha_1 Jx_n + \alpha_2 z_n + \alpha_3 T_n u_n, \\ C_{n+1} = \{z \in C_n : \|z - y_n\| \leq \|z - x_n\|\}, \\ x_{n+1} = P_{C_{n+1}} x, \end{cases}$$

for all $n \in \mathbb{N}$, $\alpha_1, \alpha_2, \alpha_3 \in (0, 1)$ such that $\alpha_1 + \alpha_2 + \alpha_3 = 1$, $\{r_n\} \subset [a, \infty)$ for some $a > 0$, $A_n = A_{n \pmod N}$ and $f_n(\cdot, \cdot) = f_{n \pmod L}(\cdot, \cdot)$. Then, $\{x_n\}$ converges strongly to $P_B x$, where P_B is the metric projection of H onto B .

Proof. In a Hilbert space, J is the identity operator and $\phi(x, y) = \|x - y\|^2$ for all $x, y \in H$. The result follows from Theorem 3.1. □

Example 4.1. [5] Let $E = l_p, 1 < p < \infty, \frac{1}{p} + \frac{1}{q} = 1$, and $C = \overline{B_{l_p}}(0, 1) = \{x \in l_p : \|x\|_{l_p} \leq 1\}$. Then $JC = \overline{B_{l_q}}(0, 1)$. Let $f : JC \times JC \rightarrow \mathbb{R}$ defined by $f(x^*, y^*) = \langle J^{-1}x^*, x^* - y^* \rangle \forall x^* \in JC, A : C \rightarrow l_q$ defined by $Ax = J(x_1, x_2, x_3, \dots) \forall x = (x_1, x_2, x_3, \dots) \in C, T : C \rightarrow l_q$ defined by $Tx = J(0, x_1, x_2, x_3, \dots) \forall x = (x_1, x_2, x_3, \dots) \in C$, and $T_n : C \rightarrow l_q$ defined by $T_n x = \alpha_n Jx + (1 - \alpha_n)Tx, \forall n \geq 1, \forall x \in C, \alpha_n \in (0, 1)$ such that $1 - \alpha_n \geq \frac{1}{2}$. Then C, JC, f, A, T , and T_n satisfy the conditions of Corollary 4.1. Moreover, $0 \in F_J(\Gamma) \cap EP(f) \cap VI(C, A)$.

5. CONCLUSION

Our theorem and its applications complement, generalize, and extend results of Uba et al. [23], Zegeye and Shahzad [27], Kumam [14], Qin and Su [19], and Nakajo and Takahashi [20]. Theorem 3.1 is a complementary analogue and extension of Theorem 3.2 of [27] in the following sense: while Theorem 3.2 of [27] is proved for a finite family of *self-maps* in uniformly smooth and strictly convex real Banach space which has the Kadec–Klee property, Theorem 3.1 is proved for countable family of *non-self maps* in uniformly smooth and uniformly convex real Banach space; in Hilbert spaces, Corollary 4.3 is an extension of Corollary 3.5 of [27] from *finite family of nonexpansive self-maps* to *countable family of nonexpansive non-self maps*. Additionally, Theorem 3.1 extends and generalizes Theorem 3.7 of [23] in the following sense: while Theorem 3.7 of [23] studied equilibrium problem and countable family of *generalized J_* -nonexpansive non-self maps*, Theorem 3.1 studied finite family of equilibrium and variational inequality problems and countable family of *generalizes J_* -nonexpansive non-self maps*; corollary 4.2 generalized Theorem 3.7 of [23] to a

finite family of equilibrium problems and countable family of *generalized J_* -nonexpansive non-self maps*. Furthermore, Corollary 4.1 extends Theorem 3.1 of [14] from Hilbert spaces to a more general uniformly smooth and uniformly convex Banach spaces and to a more general class of continuous monotone mappings. Finally, Corollary 4.1 improves and extends the results in [19, 20] from a *nonexpansive self-map* to a *generalized J_* -nonexpansive non-self map*.

6. ACKNOWLEDGEMENT

The research of the third author is supported by the Slovak Research and Development Agency under the project APVV-20-0311.

REFERENCES

- [1] Alber, Y. Metric and generalized projection operators in Banach spaces: properties and applications. In *Theory and Applications of Nonlinear Operators of Accretive and Monotone Type* (A. G. Kartsatos, Ed.), Marcel Dekker, New York (1996), pp. 15-50
- [2] Alber, Y.; and Ryazantseva, I. *Nonlinear Ill Posed Problems of Monotone Type*, Springer, London, UK, 2006
- [3] Blum, E.; and Oettli, W. From optimization and variational inequalities to equilibrium problems, *Math. Stud.* **63** (1994) 123–145
- [4] Chidume, C. E.; and Ezea, C. G. New algorithms for approximating zeros of inverse strongly monotone maps and J -fixed points, *Fixed Point Theory Appl.* **3**(2020)
- [5] Chidume, C.E.; and Nnakwe, M.O. A new Halpern-type algorithm for a generalized mixed equilibrium problem and a countable family of generalized J -nonexpansive maps, with applications, *Carpathian J. of Math.* **34**(2)(2018), 191–198
- [6] Chidume, C.E.; and Idu, K.O. Approximation of zeros of bounded maximal monotone maps, solutions of Hammerstein integral equations and convex minimization problems, *Fixed Point Theory Appl.* **97**(2016)
- [7] Chidume, C.E.; Otubo, E.E.; Ezea, C.G.; Uba, M.O. A new monotone hybrid algorithm for a convex feasibility problem for an infinite family of nonexpansive-type maps, with applications. *Adv. Fixed Point Theory* **7**(3)(2017), 413–431
- [8] Combettes, P. L.; Hirstoaga, S. A. Equilibrium programming in Hilbert spaces, *J. Nonlinear Convex Anal.* **6**(2005), 117–136
- [9] Cioranescu, I. *Geometry of Banach Spaces, Duality Mappings and Nonlinear Problems*, vol. 62, Kluwer Academic Publishers, 1990
- [10] Ibaraki, T.; Takahashi, W. A new projection and convergence theorems for the projections in Banach spaces, *J. Approx. Theory* **149**(2007), 1-14
- [11] Takahashi, W. *Nonlinear functional analysis, Fixed point theory and its applications*, Yokohama Publ., Yokohama, (2000)
- [12] Kamimura, S.; Takahashi, W. Strong convergence of a proximal-type algorithm in a Banach space, *SIAM J. Optim.* **13**(3)(2002), 938-945
- [13] Liu, B. Fixed point of strong duality pseudocontractive mappings and applications, *Abstract Appl. Anal.* **2012**, Article ID 623625, 7 pages
- [14] Kumam, P. A hybrid approximation method for equilibrium and fixed point problems for a monotone mapping and a nonexpansive, *Nonlinear Anal. Hybrid Syst.* **2**(4)(2008), 1245-1255
- [15] Nakajo, K.; Shimoji, K.; Takahashi, W. Strong convergence theorems to common fixed points of families of nonexpansive mappings in Banach spaces *J. Nonlinear Convex Anal.* ***** (2007) 11-34
- [16] Peng, J. W.; Yao J. C. A new hybrid-extragradient method for generalized mixed equilibrium problems, fixed point problems and variational inequality problems *Taiwanese J. Math.* **12**(2008), 1401-1432
- [17] Klin-eam, C.; Suantai, S.; Takahashi, W. Strong convergence theorems by monotone hybrid method for a family of generalized nonexpansive mappings in Banach spaces, *Taiwanese J. Math.* **16**(6)(2012), 1971-1989
- [18] Kohsaka, F.; Takahashi, W. Generalized nonexpansive retractions and a proximal-type algorithm in Banach spaces, *J. Nonlinear and Convex Anal.* **8**(2)(2007), 197–209
- [19] Qin, X.; Su, Y. Strong convergence of monotone hybrid method for fixed point iteration process, *J. Syst. Sci. and Complexity* **21** (2008), 474-482
- [20] Nakajo, K.; Takahashi, W. Strong convergence theorems for nonexpansive mappings and nonexpansive semi-groups, *J. Math. Anal. Appl.* **279**(2003), 372–379
- [21] Stampacchia, G. Formes bilineaires coercitives sur les ensembles convexes, *C.R. Acad. Sci. Paris*, **258**(1964), 4413-4416

- [22] Takahashi, W.; Zembayashi, K. A strong convergence theorem for the equilibrium problem with a bifunction defined on the dual space of a Banach space, *Fixed point theory and its applications, Yokohama Publ., Yokohama, (2008), 197-209*
- [23] Uba, M. O.; Otubo, E. E.; Onyido, M. A. A Novel Hybrid Method for Equilibrium Problem and A Countable Family of Generalized Nonexpansive-type Maps, with Applications, *Fixed Point Theory*, **22(1)**(2021), 359-376
- [24] Zegeye, H. Strong convergence theorems for maximal monotone mappings in Banach spaces, *J. Math. Anal. Appl.* **343**(2008), 663-671
- [25] Zegeye, H.; Shahzad, N. Strong convergence theorems for a solution of finite families of equilibrium and variational inequality problems, *Optimization* **63(2)**(2014), 207-223
- [26] Zegeye, H.; Ofoedu, E.U.; Shahzad, N. Convergence theorems for equilibrium problem, variational inequality problem and countably infinite relatively quasi-nonexpansive mappings, *Appl. Math. Comput.* **216**(2010), 3439-3449
- [27] Zegeye, H.; Shahzad, N. A hybrid scheme for finite families of equilibrium, variational inequality and fixed point problems, *Nonlinear Anal.* **74**(2011), 263-272
- [28] Zhang, S. Generalized mixed equilibrium problems in Banach spaces, *Appl. Math. Mech. -Engl. Ed.* **30(9)**(2009), 1105-1112
- [29] Dong, Q. L.; Deng, B. C. Strong convergence theorem by hybrid method for equilibrium problems, variational inequality problems and maximal monotone operators, *Nonlinear Anal. Hybrid Syst.* **4(4)**(2010), 689-698
- [30] Saewan, S.; Kumam, P. A new iteration process for equilibrium, variational inequality, fixed point problems, and zeros of maximal monotone operators in a Banach space, *J. Inequal. Appl.* **23**(2013)

UNIVERSITY OF NIGERIA
 DEPARTMENT OF MATHEMATICS
 NSUKKA - ONITSHA RD, 410001 NSUKKA, NIGERIA
Email address: markjoeuba@gmail.com

NORTHERN ILLINOIS UNIVERSITY
 DEPARTMENT OF MATHEMATICAL SCIENCES,
 DEKALB, IL 60115, UNITED STATES
Email address: mao0021@auburn.edu

COMENIUS UNIVERSITY IN BRATISLAVA
 FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS,
 MLYNSKÁ DOLINA F1, 842 48 BRATISLAVA, SLOVAK REPUBLIC
Email address: cyrilizuchukwu04@gmail.com

UNIVERSITY OF NIGERIA
 DEPARTMENT OF MATHEMATICS
 NSUKKA - ONITSHA RD, 410001 NSUKKA, NIGERIA
Email address: peter.nwokoro@unn.edu.ng